

Enumeration of maps with self avoiding loops and the $\mathcal{O}(\mathfrak{n})$ model on random lattices of all topologies

*G. Borot*¹, *B. Eynard*²

Institut de Physique Théorique,
CEA, IPhT, F-91191 Gif-sur-Yvette, France,
CNRS, URA 2306, F-91191 Gif-sur-Yvette, France.

Abstract

We compute the generating functions of a $\mathcal{O}(\mathfrak{n})$ model (loop gas model) on a random lattice of any topology. On the disc and the cylinder, they were already known, and here we compute all the other topologies. We find that they obey a slightly deformed version of the topological recursion valid for the 1-hermitian matrix models. The generating functions of genus g maps without boundaries are given by the symplectic invariants F^g of a spectral curve $\Sigma_{\mathcal{O}(\mathfrak{n})}$. This spectral curve was known before, and it is in general not algebraic.

¹ E-mail: gaetan.borot@cea.fr

² E-mail: bertrand.eynard@cea.fr

Contents

1	The $\mathcal{O}(n)$ model	2
1.1	Definition: loop gas on a random surface	2
1.2	Generating functions	4
1.3	Matrix model	5
1.4	Relation to conformal field theories	6
1.5	Boundary insertion operator	8
1.6	Loop equations	8
1.6.1	Derivation from the matrix integral	8
1.6.2	Derivation à la Tutte	10
1.7	Analyticity properties	12
1.8	Remark: convergent matrix integrals	13
2	The linear equation	14
2.1	Saddle point equation	14
2.2	Algebraic geometry construction	15
2.3	Change of variable	19
2.4	Special 1-cut solutions	20
2.5	Bilinear form	21
2.6	General 1-cut solutions	23
3	Unstable generating functions ($2 - 2g - k \geq 0$)	23
3.1	$W_1^{(0)}$ (rooted disks)	23
3.2	$y(x)$ (the spectral curve)	24
3.3	$W_2^{(0)}$ (rooted cylinders), first method	25
3.4	$W_2^{(0)}$, infinite series representation	26
3.4.1	Decomposition and characterization	26
3.4.2	Construction of a solution	27
3.4.3	Comment	27
4	Stable generating functions ($2 - 2g - k < 0$)	28
4.1	Cauchy residue formula	28
4.2	$W_k^{(g)}$ (all topologies)	29
4.2.1	Recursive residue formula for $2g - 2 + k > 0$	29
4.2.2	Comment	31
4.2.3	Proof	32
4.3	Examples	34
4.3.1	Recursion kernel at the branchpoints	34

4.3.2	$W_3^{(0)}$ (pairs of pants)	35
4.3.3	$W_1^{(1)}$ (rooted disk with one handle)	35
5	Properties of the $W_k^{(g)}$'s	36
5.1	Symmetry	37
5.2	Homogeneity	37
5.3	Special geometry	37
5.3.1	Variations of the t_j 's	38
5.3.2	Variation of t	38
5.4	Integration formula	40
5.5	F_0 (spherical maps)	41
5.5.1	Earlier results	41
5.5.2	New results	42
5.6	F_1 (toroidal maps)	43
6	Large maps	43
6.1	Principles	44
6.1.1	Asymptotic of maps	44
6.1.2	Critical points of the $\mathcal{O}(\mathfrak{n})$ model	44
6.1.3	Taking the limit $a \rightarrow 0$	45
6.2	Spectral curve in the scaling limit	47
6.3	Topological recursion in the scaling limit	49
6.4	Critical exponents	49
6.5	Remark: double scaling limit	52
7	Conclusion	52
A	Proof of the 1-cut lemma	55
B	Elliptic functions	57
C	Properties of the μ-deformed Weierstraß function	59
D	Properties of the special solutions of Eqn. 2-4	62
E	Resolvent at $a = 0$ and $b = \infty$	65

Introduction

The problem consists in counting random discrete surface, carrying random, self-avoiding, non intersecting loops, which can have $\mathbf{n}$ possible colors. In statistical physics, this is called the $\mathcal{O}(\mathbf{n})$ model (or *loop gas model*) on a random lattice, and it plays a very important role. The $\mathcal{O}(\mathbf{n})$ model on a regular lattice is one of the exactly solvable models of Baxter [2]. Its limit $\mathbf{n} \rightarrow 0$ (or more generally $\mathbf{n} = \mathbf{n}_0 + \delta\mathbf{n}$ with $\delta\mathbf{n} \rightarrow 0$) counts configurations of self-avoiding polymers in two dimensions [12]. On the random lattice, it is one of the basic toy models to understand the random geometry of discrete maps carrying structure. The $\mathcal{O}(1)$ model on random triangulations is dual to the Ising model on a random triangulations. The fully packed case of the $\mathcal{O}(\mathbf{n})$ model is dual to the $\mathbf{q}$ -Potts model on the dual of random triangulations, with $\mathbf{q} = \mathbf{n}^2$. The $\mathcal{O}(\mathbf{n})$ model with $\mathbf{n} = 2 \cos(\pi/(m+1))$ correspond the RSOS models with m states. In general, the continuum limit of the model is related to the (p, q) minimal models when $\mathbf{n} = 2 \cos(\pi p/q)$.

Matrix integrals provide powerful techniques for the combinatorics of maps [15]. The problem of counting random discrete surfaces without loops can be rephrased as a 1-matrix integral [7, 14]. Techniques to solve this 1-matrix model beyond spherical limit first appeared in [1], and culminated in the full solution in [16], by a "topological recursion" formula. This structure was later enhanced [20] to multi-matrix models, and it was used beyond matrix models to associate "symplectic invariants" to an arbitrary spectral curve.

The $\mathcal{O}(\mathbf{n})$ model was also rewritten as a matrix integral [24, 30]. In [32], then [23], the phase diagram with respect to μ such that $\mathbf{n} = -2 \cos(\pi\mu)$ was established. Besides, the critical exponents for the geometry of large maps with the topology of a disc were found. Later, a closed set of loop equations for the generating function of the $\mathcal{O}(\mathbf{n})$ maps was found [14, 18], and they were solved at least for maps with the topology of a disc or a cylinder. Some sparse other cases were investigated before [29, 31]. So far, an efficient algorithm was lacking to compute the generating functions of the $\mathcal{O}(\mathbf{n})$ model in higher topologies. It was not clear to which extent the method of [16] could be generalized. Indeed, the disc amplitude $W_1^{(0)}(x)$, which ought to give the spectral curve, is not algebraic when μ is not rational, and the loop equations seem rather different from the 1-matrix model case.

In other words, do some symplectic invariants of [20] give the solution of the $\mathcal{O}(\mathbf{n})$ model ? The answer is yes, with a slight deformation (of parameter $\mathbf{n}$) of the notion of spectral curve. Thus, we obtain all correlation functions of the random lattice for any topology and any number of boundaries.

Outline

Apart from the presentation of new explicit results for the all genus solution, our work comes in continuity with the foundation articles [30, 18, 19] on the $\mathcal{O}(\mathbf{n})$ model on random lattice despite the time gap. We also wish to make these techniques accessible to combinatorialists, who are increasingly interested in the $\mathcal{O}(\mathbf{n})$ model and the Potts model. For these reasons, we take space:

- To introduce the model and its matrix integral representation (Section 1).
- To give two derivations of the loop equations, one straightforward from the matrix integral, and the other being its combinatorial analog (Section 1.6).
- To review the solution of the master loop equation (Section 2 and 3), which was first determined in [23] for $n = 2 \cos(\pi p/q)$, in [18] for $|\mathbf{n}| \leq 2$, in [19] for $|\mathbf{n}| > 2$. Our presentation is maybe more "algebro-geometric" minded, and this step is essential for our solution to the full set of loop equations (Section 4).

Secondly, we extend to the standard properties of the topological recursion to the spectral curves of the $\mathcal{O}(\mathbf{n})$ model (Section 5). Eventually, we show how our approach allow to recover earlier results for the limit of large maps [32], and we complete them for all topologies in the light of the double scaling limit and topological recursion (Section 6).

1 The $\mathcal{O}(\mathbf{n})$ model

1.1 Definition: loop gas on a random surface

Roughly speaking, a random discrete surface³ (also called "map" in combinatorics) is a graph drawn on an oriented connected surface, such that all faces are polygons. It is thus obtained by gluing polygons along their edges. In the $\mathcal{O}(\mathbf{n})$ model, we have at our disposal empty polygons of size $j \geq 3$ (weight t_j for each), and triangles carrying a piece of path (weight $-c$ for each). A configuration correspond to a map of genus g , with v vertices, to which we give a Boltzmann weight:

$$t^v N^{2-2g} \frac{(-c)^\ell \mathbf{n}^{\#\text{loops}}}{\#\text{Aut}} \hat{t}_3^{n_3} \hat{t}_4^{n_4} \dots \hat{t}_{d_{\max}}^{n_{d_{\max}}} \quad (1-1)$$

This weight is non local because $\mathbf{n}$ is coupled to the number of loops (i.e. closed paths). The fully packed model is an interesting special case : if we set $\hat{t}_j = 0$ ($j \geq 3$), all unmarked polygons are triangles carrying a piece of path.

³See [4] for a precise definition




Figure 1: Example of a map in the $\mathcal{O}(\mathbf{n})$ model: $\mathcal{M} \in \mathbb{M}_1^{(2)}$.

In addition, we may consider maps with k marked faces, with a marked edge on each marked face. We allow marked faces having more than one side, and we call them "boundaries". For $k = 1$, one also says that the maps are "rooted".

Definition 1.1 $\mathbb{M}_k^{(g)}(v)$ is the set of connected oriented discrete surfaces of genus g , with v vertices, obtained by gluing unmarked j -gons of degree $3 \leq j \leq d_{\max}$, k marked j -gons (of degree $1 \leq j$), and triangles carrying a piece of path, such that all the paths are loops (they are automatically self-avoiding).

Proposition 1.1 $\mathbb{M}_k^{(g)}(v)$ is a finite set.

Indeed, in a configuration where marked faces have lengths $l_1, \dots, l_k$:

$$\#\text{faces} = \sum_{j=3}^{d_{\max}} n_j + k + \ell$$

The total number of edges is half the number of half-edges i.e:

$$\#\text{edges} = \frac{1}{2} \left(\sum_{j=3}^{d_{\max}} j n_j + \sum_{i=1}^k l_i + 3\ell \right)$$

The Euler characteristics is:

$$\chi = 2 - 2g = v - \#\text{edges} + \#\text{faces}$$

Hence:

$$2g - 2 + k + v = \frac{1}{2} \left(\sum_{j=3}^{d_{\max}} (j-2)n_j + \sum_{i=1}^k l_i + \ell \right) \quad (1-2)$$

When g, k, v are fixed, n_j, l_i and ℓ are bounded, and there exists only a finite number of such surfaces.

1.2 Generating functions

As a convention, $\mathbb{M}_1^{(0)}(1)$ has one element, which is a map reduced to 1 vertex.

Definition 1.2 *We define a formal series in powers of t , counting genus g maps with k marked faces:*

$$W_k^{(g)} = \sum_{v=1}^{\infty} t^v \sum_{\mathcal{M} \in \mathbb{M}_k^{(g)}(v)} \frac{(-c)^{\ell(\mathcal{M})} \mathbf{n}^{\#\text{loops}(\mathcal{M})}}{\#\text{Aut}(\mathcal{M})} \frac{\hat{t}_3^{n_3(\mathcal{M})} \hat{t}_4^{n_4(\mathcal{M})} \dots \hat{t}_{d_{\max}}^{n_{d_{\max}}(\mathcal{M})}}{(x_1 - \frac{c}{2})^{l_1(\mathcal{M})+1} \dots (x_k - \frac{c}{2})^{l_k(\mathcal{M})+1}}$$

where the i -th marked face is a $l_i(\mathcal{M})$ -gon. For closed maps, we call $W_0^{(g)} = F_g$.

This makes sense because the coefficient of t^v is a finite sum. It is in fact polynomial in $\mathbf{n}$ and $\hat{t}_j$'s, and a rational fraction in x_i 's with poles only at $x_i = \frac{c}{2}$. In particular for rooted maps, $\#\text{Aut}(\mathcal{M}) = 1$. For instance:

$$W_1^{(0)} = \frac{t}{x - \frac{c}{2}} + \sum_{v \geq 2} t^v \sum_{\mathcal{M} \in \mathbb{M}_1^{(0)}(v)} (-c)^{\ell(\mathcal{M})} \mathbf{n}^{\#\text{loops}} \frac{\hat{t}_3^{n_3(\mathcal{M})} \hat{t}_4^{n_4(\mathcal{M})} \dots \hat{t}_{d_{\max}}^{n_{d_{\max}}(\mathcal{M})}}{(x - \frac{c}{2})^{l_1(\mathcal{M})+1}}$$

Definition 1.3 *We may collect all genera:*

$$\ln Z = \sum_{g=0}^{\infty} \left(\frac{N}{t} \right)^{2-2g} F_g, \quad W_k = \sum_{g=0}^{\infty} \left(\frac{N}{t} \right)^{2-2g-k} W_k^{(g)}$$

Like in [7], all the relations between generating functions which appear in this article must be understood as equalities between formal power series of t . To each power t^v , the sum over g in the right hand side ranges over a finite number of maps, with a maximal genus $g_{\max}(v)$. There is no problem of exchange of limits. Notice that, according to Eqn. 1-2, $2g - 2 + k + v \geq 0$, so that only nonnegative powers of t appear in W_k . Z is by construction the generating function of $\mathcal{O}(\mathbf{n})$ maps (possibly not connected) with weight given by Eqn. 1-1.

1.3 Matrix model

[7] opened the way to represent these partition functions as formal matrix integrals. The $\mathcal{O}(\mathfrak{n})$ matrix model was first introduced by I. Kostov [30]:

$$Z = \int_{\text{formal}} d\hat{M} dA_1 \cdots dA_n e^{-\frac{N}{t} \text{Tr} \left[\hat{V}(\hat{M}) + (\hat{M} + \frac{c}{2}) \sum_i A_i^2 \right]} \quad (1-3)$$

$d\hat{M}$ and dA_i are the usual Lebesgue measures on the vector space $\mathcal{H}_N$ of $N \times N$ hermitian matrices. $V(\hat{M})$ is the potential:

$$\hat{V}(\hat{M}) = \frac{\hat{M}^2}{2} - \sum_{j=3}^{d_{\max}} \frac{\hat{t}_j \hat{M}^j}{j} \equiv \frac{\hat{M}^2}{2} - \hat{V}_{\geq 3}(\hat{M})$$

The notation $\int_{\text{formal}}$ actually means:

$$Z = \frac{1}{Z_0} \sum_{j=0}^{\infty} \frac{N^j t^{-j}}{j!} \int_{\mathbf{H}_N^{n+1}} d\hat{M} dA_1 \cdots dA_n e^{-\frac{N}{2t} \text{Tr} \left(\hat{M}^2 + c \sum_{i=1}^n A_i^2 \right)} \left[\text{Tr} \left(\hat{V}_{\geq 3}(\hat{M}) - \sum_{i=1}^n \hat{M} A_i^2 \right) \right]^j$$

where Z_0 is the Gaussian integral:

$$Z_0 \equiv \int_{\mathbf{H}_N^{n+1}} d\hat{M} dA_1 \cdots dA_n e^{-\frac{N}{2t} \text{Tr} \left(\hat{M}^2 + c \sum_{i=1}^n A_i^2 \right)} = 2^{\frac{(n+1)N}{2}} c^{-\frac{nN^2}{2}} \left(\frac{\pi t}{N} \right)^{\frac{(n+1)N^2}{2}}$$

The coefficient of t^v is a finite sum of moments of gaussian integrals over hermitian matrices of size $N \times N$. This model is defined for any $\mathfrak{n} \in \mathbb{C}$. Z , $\ln Z$, $\dots$ are well defined formal power series in t , and they coincide [24] with the generating functions of maps in the $\mathcal{O}(\mathfrak{n})$ model. For instance:

$$W_k = \sum_{g=0}^{\infty} \left(\frac{N}{t} \right)^{2-2g-k} W_k^{(g)} = \left\langle \prod_{i=1}^k \text{Tr} \frac{1}{x_i - \frac{c}{2} - \hat{M}} \right\rangle_{C, \text{formal}}$$

C stands for "cumulant", the expectation value is taken with respect to the formal measure of Eqn. 1-3 and:

$$\left(\frac{1}{x_i - \frac{c}{2} - \hat{M}} \right)_{\text{formal}} \equiv \sum_{j=0}^{\infty} \frac{\hat{M}^j}{(x_i - \frac{c}{2})^{j+1}}$$

It is more convenient to study the following matrix model, where we shifted to $M = \hat{M} + \frac{c}{2}$:

$$Z = \int_{\text{formal}} dM dA_1 \cdots dA_n e^{-\frac{N}{t} \text{Tr} \left(V(M) + \sum_{i=1}^n M A_i^2 \right)} \quad (1-4)$$

$$W_k = \left\langle \prod_{i=1}^k \text{Tr} \frac{1}{x_i - M} \right\rangle_C \quad (1-5)$$

where the expectation value is taken with respect to the measure of Eqn. 1-4, and the potential is:

$$V(M) = \hat{V}(\hat{M}) \equiv t_0 + \sum_{j=1}^{d_{\max}} \frac{t_j M^j}{j}$$

To recover combinatorial quantities, one has to expand the correlation functions first as formal series in t , then as Laurent series in $\hat{x}_i = x_i - \frac{c}{2}$. W_k and $W_k^{(g)}$ are called correlation functions in the language of statistical physics. The fully packed case correspond to the quadratic potential

$$V_{\text{FPL}}(x) = \frac{1}{2} \left(x - \frac{c}{2} \right)^2$$

The A matrices are gaussian and can be integrated out. Then, $W_k^{(g)}$'s appear as correlators of eigenvalues (λ_i) of M , with a weight on eigenvalues:

$$p(\lambda_1, \dots, \lambda_N) \propto \frac{\prod_{1 \leq i < j \leq N} (\lambda_i - \lambda_j)^2}{\left[\prod_{1 \leq i, j \leq N} (\lambda_i + \lambda_j) \right]^{n/2}}$$

1.4 Relation to conformal field theories

Polygonal large maps constructed by matrix models provide a discretization of Riemann surfaces. In the continuum limit, physically speaking, it is thought to define a theory of 2D quantum gravity: observables should be weighted sums over all possible surfaces endowed with a metric. The self avoiding paths present in the $\mathcal{O}(n)$ model are in some sense matter fields added on the surface. The usual approach of 2D quantum gravity is Liouville field theory, which is a statistical model on the moduli space of Riemann surfaces endowed with a metric (see [38] for a review). Liouville theory can also be coupled to matter fields. Both approaches are conjectured to coincide and to enjoy conformal invariance.

"Double scaling limits" of matrix models are conjectured⁴ to be conformal field theories (CFT) coupled to gravity [11]. This means in practice that the scaling exponents are given by Kac's table, that the double scaling limit (the definition is recalled in Section 6.5) W_k^* of the correlation functions satisfy PDE's on the spectral curve. These equations come from a representation of a Virasoro algebra with central charge c . In this correspondence, the double scaling limit of the $\mathcal{O}(n)$ matrix model should be a conformal theory with central charge:

$$c = 1 - 6 \left(\sqrt{g} - \frac{1}{\sqrt{g}} \right)^2 \quad (1-6)$$

⁴To our knowledge, this conjecture is up to now proved only for the 1-matrix model, near an edge point where the equilibrium density of eigenvalues $y(x)$ behaves as $x^{p/2}$ [5]

where $\mathbf{g}$ is such that $\mathbf{n} = -2 \cos(\pi \mathbf{g})$. Various $\mathbf{g}$ corresponding to the same $\mathbf{n}$ define the various *phases* of the $\mathcal{O}(\mathbf{n})$ model. Many studies on the $\mathcal{O}(\mathbf{n})$ model with boundary operators [35, 34, 6, 32] support this proposal. See also the review [10] and the article [37].

We bound ourselves to underline that, if those conjectures were correct, the critical exponents of the $\mathcal{O}(\mathbf{n})$ model would be known for any topology from the KPZ relation [27, 8, 9, 38]. KPZ is derived from Liouville conformal field theory and expresses how the critical exponents change when a CFT is coupled to gravity. Independently of CFT, rigorous results for the $\mathcal{O}(\mathbf{n})$ model were derived by sheer analysis of the loop equations, and they agree with the CFT predictions. In the literature, the exponents and the scaling form of the following *genus 0* functions of Fig. 2 were known.




Figure 2:

In this article, we obtain (Section 6) the exponents and the scaling forms in all topologies of functions without loop insertion on the boundaries (Fig. 3).




Figure 3:

1.5 Boundary insertion operator

We can mark a face of size j , by taking a derivative with respect to t_j :

$$W_{k+1}^{(g)}(x_1, \dots, x_k, x_{k+1}) = \sum_{j \geq 0} \frac{1}{(x_{k+1} - \frac{c}{2})^{j+1}} \frac{\partial W_k^{(g)}(x_1, \dots, x_k)}{\partial \hat{t}_j}$$

Although we defined $\hat{t}_j$ only for $j \geq 3$ (this is the condition for Prop. 1.1 to hold), it is allowed to define $\hat{t}_2$, $\hat{t}_1$ and $\hat{t}_0$, and derivatives $\partial/\partial \hat{t}_0$ at $\hat{t}_0 = 0$, $\partial/\partial \hat{t}_1$ at $\hat{t}_1 = 0$, and $\partial/\partial \hat{t}_2$ at $\hat{t}_2 = -1$. So, we define the "boundary insertion operator" as the formal series:

$$\frac{\partial}{\partial \hat{V}(\hat{x})} \stackrel{\text{def}}{=} \sum_{j \geq 0} \frac{j}{\hat{x}^{j+1}} \frac{\partial}{\partial \hat{t}_j}$$

In the new variable, $x = \hat{x} + \frac{c}{2}$, which is a formal series of the former one, we can do the resummation:

$$\frac{\partial}{\partial \hat{V}(\hat{x})} = \sum_{j \geq 0} \frac{j}{x^{j+1}} \frac{\partial}{\partial t_j} \equiv \frac{\partial}{\partial V(x)}$$

The reason for this definition is:

$$W_{k+1}^{(g)}(x_1, \dots, x_k, x_{k+1}) = \frac{\partial}{\partial V(x_{k+1})} W_k^{(g)}(x_1, \dots, x_k) \quad (1-7)$$

1.6 Loop equations

The loop equations provide relationships between the generating functions $W_k^{(g)}$'s. They can be derived either by integration by parts in the matrix integral, or by combinatorial manipulations. The first method is much faster, but the reader not familiar with matrix integrals may prefer the bijective proof. For completeness we recall the two possibilities.

1.6.1 Derivation from the matrix integral

Loop equations are the infinitesimal counterparts of the invariance of an integral under a continuous family of change of variable [36]. They are true both for formal integrals and convergent integrals, so we do not have to bother with variable $\hat{x}$, and we can work directly with variable x . One has to compute the jacobian of the change of variable and the variation of the exponential term. The loop equation merely states that the jacobian cancels the variation of the exponential, in expectation. The general method for computing jacobians of infinitesimal changes of matrix variables, is called "split and merge", and is exposed in many articles (for instance, [15]).

Let us write $I = \{x_2, \dots, x_k\}$, and define the following auxiliary functions:

$$G_k = \sum_{g=0}^{\infty} \left(\frac{N}{t} \right)^{2-2g-k} G_k^{(g)} = \left\langle \text{Tr} \frac{1}{x-M} A_1^2 \prod_{x_i \in I} \text{Tr} \frac{1}{x_i-M} \right\rangle_C$$

$$\begin{aligned}\tilde{G}_k &= \sum_{g=0}^{\infty} \left(\frac{N}{t}\right)^{2-2g-k} \tilde{G}_k^{(g)} = \left\langle \text{Tr} \frac{1}{x-M} A_1 \frac{1}{x'-M} A_1 \prod_{x_i \in I} \text{Tr} \frac{1}{x_i-M} \right\rangle_C \\ P_k &= \sum_{g=0}^{\infty} \left(\frac{N}{t}\right)^{2-2g-k} P_k^{(g)} = \left\langle \text{Tr} \frac{V'(x) - V'(M)}{x-M} \prod_{x_i \in I} \text{Tr} \frac{1}{x_i-M} \right\rangle_C\end{aligned}$$

$P_k^{(g)}$ is by construction the polynomial part of $V'(x) W_k^{(g)}(x, x_2, \dots, x_k)$ at large x :

$$P_k^{(g)}(x, x_2, \dots, x_k) = \left(V'(x) W_k^{(g)}(x, x_2, \dots, x_k) \right)_+$$

We indicate infinitesimal change of variable and the corresponding loop equations:

- With $M \rightarrow M + \epsilon \frac{1}{x-M}$, we obtain to first order in ϵ :

$$W_1(x)^2 + W_2(x, x) = \frac{N}{t} \left[V'(x) W_1(x) - P_1(x) - n G_1(x) \right] \quad (1-8)$$

- With $A_1 \rightarrow A_1 + \epsilon \frac{1}{x-M} A_1 \frac{1}{x'-M}$, we obtain:

$$W_1(x) W_1(x') + W_2(x, x') = \frac{N}{t} \left[(x + x') \tilde{G}_1(x, x') - G_1(x) - G_1(x') \right] \quad (1-9)$$

Auxiliary functions can be eliminated by specializing to $x = -x'$ and combining the two equations:

$$\begin{aligned} & W_2(x, x) + n W_2(x, -x) + W_2(-x, -x) + W_1(x)^2 + n W_1(x) W_1(-x) + W_1(-x)^2 \\ &= \frac{N}{t} [V'(x) W_1(x) + V'(-x) W_1(x) - P_1(x) - P_1(-x)]\end{aligned}$$

If we collect the highest power in N , which is N^2 , we obtain:

Theorem 1.1 *Master loop equation (quadratic in $W_1^{(0)}$).*

$$\begin{aligned} & W_1^{(0)}(x)^2 + n W_1^{(0)}(x) W_1^{(0)}(-x) + W_1^{(0)}(-x)^2 \\ &= V'(x) W_1^{(0)}(x) + V'(-x) W_1^{(0)}(x) - P_1^{(0)}(x) - P_1^{(0)}(-x)\end{aligned}$$

We stress that Eqns. 1-8 and 1-9 are true for any potential for which the quantities involved make sense. In particular, (t_0, t_1, t_2) is not restricted to $(0, 0, -1)$. Accordingly, we can obtain loop equations for all W_k by successive applications of $\frac{\partial}{\partial V}$. Here is the general results for $k \geq 1$ and collecting the N^{2-2g} terms ($g \geq 0$).

Theorem 1.2 *Higher loop equations (linear in $W_k^{(g)}$).*

$$\begin{aligned} & W_{k+1}^{(g-1)}(x, x, I) + W_{k+1}^{(g-1)}(-x, -x, I) + n W_{k+1}^{(g-1)}(x, -x, I) \\ &+ \sum_{J \subseteq I, 0 \leq h \leq g} \left(W_{|J|+1}^{(h)}(x, J) W_{k-|J|}^{(g-h)}(x, {}^c J) + W_{|J|+1}^{(h)}(-x, J) W_{k-|J|}^{(g-h)}(-x, {}^c J) \right. \\ &\quad \left. + n W_{|J|+1}^{(h)}(x, J) W_{k-|J|}^{(g-h)}(-x, {}^c J) \right) \\ &+ \sum_{x_i \in I} \frac{\partial}{\partial x_i} \left(\frac{W_{k-1}^{(g)}(x, I \setminus \{x_i\}) - W_{k-1}^{(g)}(I)}{x - x_i} - \frac{W_{k-1}^{(g)}(-x, I \setminus \{x_i\}) - W_{k-1}^{(g)}(I)}{x + x_i} \right) \\ &= V'(x) W_k^{(g)}(x, I) + V'(-x) W_k^{(g)}(-x, I) - P_k^{(g)}(x, I) - P_k^{(g)}(-x, I)\end{aligned}$$


We take $W_k^{(-1)} = 0$ as convention. $\sum'$ means that we exclude the two terms where W_k^g itself appears.

Thm. 1.1 is just a specialization of Thm. 1.2 to $(k, g) = (1, 0)$, but the master loop equations plays a special role. When $\mathbf{n} = 0$, we recover the loop equations of maps [40, 41, 15].

1.6.2 Derivation à la Tutte


Now, we give a bijective proof of Thm. 1.1. We introduce the auxiliary functions:

- $\tilde{G}_k^{(g)}(x, x', I)$, which counts maps which carry a path of weight 1 (and not $\mathbf{n}$) starting on a distinguished marked face (of length $l + l' + 2$), and ending on the same marked face. l is the notation for the distance between the starting point and ending point, and the weight is $\propto x^{-(l+1)} x'^{-(l'+1)}$.




- $G_k^{(g)}(x, I)$, defined by the same sum restricted to maps where the starting point is just next to the ending point ($l \equiv 1$). The weight is $\propto x'^{-(l'+1)}$. Actually:

$$G_k^{(g)}(x') = \lim_{x \rightarrow \infty} x \tilde{G}_k^{(g)}(x, x')$$



The combinatorial derivation follows the method of Tutte [40, 41], i.e we look for a recursion on the number of edges.

- Consider $W_1^{(0)}$, which counts rooted planar maps. If we remove the marked edge in a configuration counted by $W_1^{(0)}$, we count $\left[\left(x - \frac{c}{2} \right) W_1^{(0)}(x) - t \right]$ (taking care of the planar map with only 1 vertex) $W_1^{(0)}$. Three cases occur when we remove the marked edge. the face on the other side of the marked edge could be an unmarked polygon, it could carry a piece of path, or it could be the marked face itself. Pictorially, if we fill the marked face in dark grey (the whole map should be a sphere): Thus:



$$\left(x - \frac{c}{2} \right) W_1^{(0)}(x) - t = \left[\hat{V}_{\geq 3} \left(x - \frac{c}{2} \right) W_1^{(0)}(x) \right]_- + n G_1^{(0)}(x) + W_1^{(0)}(x)^2$$

where $(\cdots)_-$ means the negative part of the Laurent expansion in $x - \frac{c}{2}$. We see that the whole $\hat{V}'(x - \frac{c}{2}) = V'(x)$ is restored in this equation: nothing special


happens with the quadratic part, and the loop equation takes a nice form because of the convention for the term $v = 1$ in $W_1^{(0)}$. The polynomial (in the variable $\hat{x}$ or x):

$$P_1^{(0)}(x) = \left\langle \frac{\hat{V}'(\hat{x}) - \hat{V}'(\hat{M})}{\hat{x} - \hat{M}} \right\rangle^{(0)} = \left\langle \frac{V'(x) - V'(M)}{x - M} \right\rangle^{(0)}$$

is precisely the nonnegative part of the Laurent expansion of $V'(x)W_1^{(0)}(x)$. Hence Eqn. 1-8:

$$W_1^{(0)}(x)^2 = V'(x)W_1^{(0)}(x) - P_1^{(0)}(x) - nG_1^{(0)}(x)$$

- Consider configurations counted by $\tilde{G}_1^{(0)}(x, x')$. If we remove the triangle of the marked face where the path starts, we are counting $\left[(x + x' - c) \tilde{G}_1^{(0)}(x, x') - G_1^{(0)}(x) - G_1^{(0)}(x') \right]$. Indeed, either l or l' is shortened by one, but we have to take care of the degenerate cases $G_1^{(0)}(x)$ and $G_1^{(0)}(x')$ where l' (resp. l) shrinks to 0. There are two possibilities for this removal: either the path was of length 0 or not. Thus, we find Eqn. 1-9:



$$((x + x' - c) \tilde{G}_1^{(0)}(x, x') - G_1^{(0)}(x) - G_1^{(0)}(x')) = W_1^{(0)}(x)W_1^{(0)}(x') - c \tilde{G}_1^{(0)}(x, x')$$

Higher loop equations (Thm. 1.2) can be derived in a similar way.

1.7 Analyticity properties

To each order t^v , the sum over $M_k^{(g)}(v)$ in $W_k^{(g)}$ is a finite sum, and in particular, the coefficient of t^v in $W_k^{(g)}$ is a rational fraction of each x_i , with poles only at $x_i = \frac{c}{2}$, of maximal degree $4g - 4 + k + 2v$. In Appendix A, we prove that this implies the following analytical properties for the $W_k^{(g)}$'s:

Lemma 1.1 *1-cut lemma.*

There exists $\rho_0 > 0$ (depending only on the degree $d_{\max}$ of the polynomial $\hat{V}$, $\hat{t}_3, \dots, \hat{t}_{d_{\max}}$, $\mathbf{n}$ and c , and a priori on k, g) such that, for all $r > 0$:

- If $I = (x_1, \dots, x_k) \in (\mathbf{C} \setminus \mathcal{D}(\frac{c}{2}; r))^k \Rightarrow$, then $W_k^{(g)}(I)$, as a formal series in t , has a radius of convergence $\geq \frac{r}{\rho_0}$.

Accordingly, if we hold $k - 1$ variables $J = (x_2, \dots, x_k)$ fixed, at values different from $\frac{c}{2}$: for $r > 0$ small enough, for all $t \in \mathcal{D}(0; \frac{r}{\rho_0})$, $W_k^{(g)}(x, J)$ is holomorphic for $x \in \mathbf{C} \setminus \mathcal{D}(\frac{c}{2}; r)$, $\mathcal{C}^\infty$ in t , and $\mathcal{D}(\frac{c}{2}; r) \cap \overline{\mathcal{D}}(-\frac{c}{2}; r) = \emptyset$.

More precisely, there exists $t_0 > 0$ (depending only on $d_{\max}, \hat{t}_3, \dots, \hat{t}_{d_{\max}}, \mathbf{n}$ and c) and two formal series $a(t), b(t)$ in $\sqrt{t}$, whose radius of convergence in $\sqrt{t}$ is greater than $\sqrt{t_0}$ (non zero), such that (we write $\gamma(t)$ the segment $[a(t), b(t)]$):

- $a(t) = \frac{c}{2} - c_1\sqrt{t} + O(\sqrt{|t|})$ and $b(t) = \frac{c}{2} + c_1\sqrt{t} + O(\sqrt{|t|})$ for some $c_1 \in \mathbb{C}$.
- $W_1^{(0)}(x)$ is absolutely convergent for $t \in \mathcal{D}(0; t_0)$, holomorphic on $\mathbf{C} \setminus \gamma(t)$, and has a discontinuity on $\gamma(t)$. On a neighborhood of $\gamma(t)$, it takes the form $h(x) \times \sqrt{(x-a)(x-b)}$, where h is meromorphic in a neighborhood of $\gamma(t)$, has no pole except maybe in $a(t)$ and $b(t)$, and has no zeroes on $\gamma(t)$. We call this behavior a square root discontinuity.
- For all the other k, g 's, $W_k^{(g)}(x_1, \dots, x_k)$ has a square root discontinuity in each variable on $\gamma(t)$, and is holomorphic on $\mathbb{C} \setminus \gamma(t)$.

This lemma is quite technical, and we give its proof in Appendix A. As a brief sketch of the proof, let us say that, by a very rough bound on $\#\mathbb{M}_1^{(0)}(v)$, we first prove that $W_1^{(0)}$ is convergent in the domain $\mathbb{C} \setminus \mathcal{D}(\frac{c}{2}; \frac{|c|}{2})$ when $|t| < t^*$ for some t^* , and thus $W_1^{(0)}(x)$ can have singularities only in a small disc centered in c , of radius smaller than $\frac{|c|}{2}$. This implies that $W_1^{(0)}(-x)$ is analytical for x in this disc. Then, from the master loop equation (Thm. 1.1), $W_1^{(0)}$ can only have (and must have) square-root discontinuity in the disc, at points $x = a(t)$ and $x = b(t)$. Eventually, the series $a(t)$ and $b(t)$ are determined by the master loop equation. This property is called the 1-cut assumption in physics (although it is not an assumption here), and is closely related to Brown's lemma in combinatorics. The loop equations themselves do not have a unique solution, but we shall see that there is a unique one satisfying this lemma.

1.8 Remark: convergent matrix integrals

The 1-cut assumption holds for formal matrix integrals, i.e generating functions of the $O(\mathbf{n})$ model configurations.

However, one could be interested in studying the matrix integral of Eqn. 1-4, not as a formal matrix integral, but as a genuine convergent integral. In this case, a 1-cut lemma can hold or not, depending on the choice of $V(M)$, and in fact on the choice of the integration domain for the eigenvalues of M . In some sense, it holds if the integration domain is a "steepest descent" integration path for the potential V , but those considerations are beyond the scope of our article. When this is the case, it is known that, for $|t|$ small enough according to the bound of the 1-cut lemma, W_k does have an asymptotic expansion of the form:

$$W_k^{(g)} = \sum_{g=0}^{\infty} \left(\frac{N}{t} \right)^{2-2g-k} W_k^{(g)}(t) \quad (1-10)$$

Of course, these $W_k^{(g)}$ satisfy then the loop equations (Thm 1.2). In this article, we shall consider only the situation (realized in combinatorics, because of Lemma 1.1) where the 1-cut property holds, with the cut $[a, b] \subseteq \mathbb{R}_+$. We leave the "multi-cut solution" to $\mathcal{O}(\mathfrak{n})$ model loop equations for a future work.

2 The linear equation

We write $\mathfrak{n} = -2 \cos(\pi\mu)$ and we assume $\mu \in]0, 1[\setminus \{0\}$. $\mu \in]0, 1[$ is in bijection with $\mathfrak{n} \in]-2, 2[$. With few technical modifications, the loop equations can also be solved for $\mathfrak{n}$ outside of that range⁵, but the nature of their solution is quite different. The range $\mathfrak{n} \in \mathbb{R}$ was analyzed in [19]. We now review the techniques developed in [14] to solve the master loop equation (Thm. 1.1), and introduce enough algebraic geometry to present the solution of the higher loop equations.

2.1 Saddle point equation

Due to the analytical structure of $W_1^{(0)}$ (square root discontinuity with end points $a(t), b(t)$), we can transform the master loop equation, which is quadratic, into a linear one. The latter was originally called "saddle point equation" because it coincides with the saddle point approximation for the density of eigenvalues [23].

Proposition 2.1

$$\forall x \in [a(t), b(t)] \quad W_1^{(0)}(x + i\epsilon) + \mathfrak{n} W_1^{(0)}(-x) + W_1^{(0)}(x - i\epsilon) \underset{\epsilon \rightarrow 0}{=} V'(x)$$

(2-1)

⁵Notice the $\mathcal{O}(-2)$ model is dual to the $\mathcal{O}(0)$ model by the change of variable $M \rightarrow \sqrt{M}$, i.e. to the 1-hermitian matrix model with a non analytic potential $\tilde{V}(M) = V(\sqrt{M})$

proof:

We start from Thm. 1.1

$$\begin{aligned} & W_1^{(0)}(x)^2 + nW_1^{(0)}(x)W_1^{(0)}(-x) + W_1^{(0)}(-x)^2 \\ = & V'(x)W_1^{(0)}(x) + V'(-x)W_1^{(0)}(-x) - P_1^{(0)}(x) - P_1^{(0)}(-x) \end{aligned}$$

where $P_1^{(0)}(x)$ is a polynomial in x of degree $(\deg V - 2)$. Because of the 1-cut lemma:

$$\forall x \in [a(t), b(t)] \quad W_1^{(0)}(-x + i\epsilon) - W_1^{(0)}(-x - i\epsilon) \underset{\epsilon \rightarrow 0}{=} 0$$

Besides, let us define:

$$\begin{aligned} & 2P_e(x) \\ = & P_1^{(0)}(x) + P_1^{(0)}(-x) \\ = & W_1^{(0)}(x)^2 + nW_1^{(0)}(x)W_1^{(0)}(-x) + W_1^{(0)}(-x)^2 - V'(x)W_1^{(0)}(x) - V'(-x)W_1^{(0)}(-x) \end{aligned}$$

$P_e(x)$ is a polynomial, so:

$$\forall x \in [a(t), b(t)] \quad P_e(x + i\epsilon) - P_e(x - i\epsilon) \underset{\epsilon \rightarrow 0}{=} 0$$

This equation factorizes into:

$$(W_1^{(0)}(x + i\epsilon) - W_1^{(0)}(x - i\epsilon)) \left[W_1^{(0)}(x + i\epsilon) + nW_1^{(0)}(-x) + W_1^{(0)}(x - i\epsilon) - V'(x) \right] \underset{\epsilon \rightarrow 0}{=} 0$$

Since on $x \in [a(t), b(t)]$ we have $W_1^{(0)}(x + i\epsilon) \neq W_1^{(0)}(x - i\epsilon)$ in the limit $\epsilon \rightarrow 0$, we find Eqn. 2-1. $\square$

Eqn. 2-1 is linear, provided that $a(t)$ and $b(t)$ are known. The non-linearity is hidden in the determination of $a(t)$ and $b(t)$. Given a segment $[a, b]$ of the positive real line, we shall study the general 1-cut solutions of the homogeneous linear equation:

$$\forall x \in [a, b] \quad W(x + i\epsilon) + nW(-x) + W(x - i\epsilon) \underset{\epsilon \rightarrow 0}{=} 0 \quad (2-2)$$

The extension to an arbitrary path $[a, b]$ in the complex plane presents no difficulty, but is not needed for combinatorics.

2.2 Algebraic geometry construction

We look for a solution of Eqn. 2-1, with only one cut $[a, b]$, and in particular which is analytical on $[-b, -a]$. Since our equation involves both $x \in [a, b]$ and $-x \in [-b, -a]$, it is convenient to introduce a conformal mapping between the complex plane with two cuts $[a, b] \cup [-b, -a]$ and the hyperelliptical curve $\mathcal{S} : \sigma^2 = (x^2 - a^2)(x^2 - b^2)$.

We assume $0 < a < b < \infty$. The cases $a = 0$ or $b = \infty$, treated directly in Section E, are easier and somewhat more explicit, and they correspond to critical points [32], as we review in Section 6.1.

+ - sign of (Re σ , Im σ)

Square root determination :

— Re $\sigma = 0$

$$\sigma^2 = \rho e^{i\theta}$$

$$\square \quad \sigma = \sqrt{\rho} e^{i\theta/2}$$

— $\sigma^2 \in \mathbf{R}_+$

$$\rho \in \mathbf{R}_+$$

$$\square \quad \sigma = -\sqrt{\rho} e^{i\theta/2}$$

— $\sigma^2 \in \mathbf{R}_-$

$$\theta \in]-\pi, \pi[$$




Figure 4: $\mathcal{S} : \sigma^2 = (x^2 - a^2)(x^2 - b^2)$ has two sheets, corresponding to opposite square roots, and we call one of them the *physical sheet*. The determination (white or yellow) of the square root changes when one crosses the lines where $\sigma^2 \in \mathbf{R}_-$, so that σ is an analytical function of x on each sheet, outside $[-b, -a] \cup [a, b]$. We have indicated the corresponding signs for $\text{Re}(\sigma)$ and $\text{Im}(\sigma)$ on the two sheets.




Figure 5:

$\mathcal{S}$ is topologically a torus. So, the space of holomorphic differential form is of complex dimension 1 and is generated by dx/σ . To parametrize the surface, we define

$$u(x) = \frac{ib}{2K'} \int_{-a}^x \frac{dx'}{\sigma(x')}$$

which is path dependant. We fix $K' \in \mathbb{R}_+^*$ such that $u(-b) = \frac{1}{2}$ along the blue path. Then, $u(a) = \tau$, shown in the next paragraph to be half of the modulus of the torus, lies in $i\mathbb{R}_+$ for $a, b \in \mathbb{R}_+$, $a < b$. We have drawn the paths followed on the first sheet only: they follow the real line, the blue one on the side $\text{Im } x > 0$, the purple one on the side $\text{Im } x < 0$. Because the square root on the second sheet is opposite to its determination on the physical sheet, the analogue integration paths on the second sheet lead to opposite u for the same value of x .




Figure 6:

We present the u -plane at the end of the construction. Circling only once along the paths in the two sheets gives the rectangular region $[-\frac{1}{2}, \frac{1}{2}] \times [-\tau, \tau]$. The path corresponding to $u \in [-\frac{1}{2}, \frac{1}{2}]$ or $u \in [-\tau, \tau]$ are non contractible loops in $\mathcal{S}$. Following these paths adds 1 (or 2τ) to u and leads to the same starting point. Then, it is Abel's theorem that u induces an isomorphism between $\mathcal{S}$ and the Riemann surface $\mathbb{C}/(\mathbb{Z} \oplus 2\tau\mathbb{Z})$. Moreover, we have marked the points corresponding to $x + i\epsilon$, $x - i\epsilon$ ($\epsilon > 0$), and $-x$ for some $x \in [a, b]$. We see that, in the variable u , they are merely τ -translations of each other:

generic point	images in the torus
x	$u, -u$
$-x$	$\tau - u, -(\tau - u)$
$\bar{x}$	$2\tau - u, u - 2\tau$

In a nutshell, our parametrization in terms of elliptic functions [25] reads:

$$x = a \operatorname{sn}_k(\varphi), \quad \varphi = \int_0^{x/a} \frac{dx}{\sqrt{(1-x'^2)(1-k^2x'^2)}}, \quad u = \frac{i\varphi}{2K'} + \frac{\tau}{2}$$

where sn_k is the odd solution of $y' = \sqrt{(1-y^2)(1-k^2y^2)}$, and:

$$\begin{aligned} \operatorname{cn}_k &= \sqrt{1 - \operatorname{sn}_k^2}, & \operatorname{dn}_k &= \sqrt{1 - k^2 \operatorname{sn}_k^2}, & k &= a/b, & \tau &= \frac{iK}{K'} \\ K &= K(k) = \int_0^1 \frac{dx'}{\sqrt{(1-x'^2)(1-k^2x'^2)}} & & \text{is the complete elliptic integral} \\ K' &= K(\sqrt{1-k^2}) = \int_0^\infty \frac{dx'}{\sqrt{(1+x'^2)(1+k^2x'^2)}} \end{aligned}$$

With help of the properties [25] under translation and rotation to imaginary argument of the elliptic functions sn , cn , dn , we obtain:

$$x = a \operatorname{dn}_{\sqrt{1-a^2/b^2}} \left(\frac{2u}{K'} \right)$$

We mention it to be explicit, though it is not at all necessary to know this formula in what follows.

2.3 Change of variable

This construction provides a nice change of variable to solve Eqn. 2-2. Let us note the inverse function of $x \mapsto u(x)$:

$$\begin{array}{ccc} x : [-\frac{1}{2}, \frac{1}{2}] \times [0, \tau] & \rightarrow & \mathbb{C}, \text{ physical sheet} \\ u & \mapsto & x(u) \end{array}$$

Then, any function:

$$\begin{array}{ccc} W : \mathbb{C} & \rightarrow & \mathbb{C} \\ x & \mapsto & W(x) \end{array}$$

which is at least analytic in the x -plane with two cuts $[-b, -a]$ and $[a, b]$, defines without ambiguity an analytic function:

$$\begin{array}{ccc} W : [-\frac{1}{2}, \frac{1}{2}] \times [0, \tau] & \rightarrow & \mathbb{C} \\ u & \mapsto & W(x(u)) \end{array}$$

In some cases, W may be extended to the whole u -plane because values on the boundary of the initial region match. Eventually, properties of $W(x)$ are translated into properties of $W(u)$ and vice versa.

in the x plane	in the u plane
no cut on $[\infty, -b] \cup [b, \infty]$	1-translation invariant
no cut on $[-b, -a]$	even
(no cut on $[-b, -a]) \times \sigma(x)$	odd
$R_e(x) + \sigma(x)R_o(x)$ R_e even rational function of x R_o odd rational function of x	1- and τ -translation invariant

Figure 7: Dictionary between invariances in variable u and analyticity properties in variable x .

Now, we apply this to any 1-cut solution W of Eqn. 2-2. We end up with a meromorphic function W defined on the whole u -plane, which is 1-translation invariant, u -even, and satisfies:

$$\forall u \in \tau + \left[-\frac{1}{2}, \frac{1}{2}\right] \quad W(2\tau - u) + \mathbf{n}W(\tau - u) + W(u) = 0 \quad (2-3)$$

Since W is analytic, it must be true on the whole u -plane. Using the parity property, and defining the operators of 1-translation $\mathbf{T}_1$, and of τ -translation $\mathbf{T}$, we rewrite Eqn. 2-3 as:

$$\forall u \in \mathbb{C} \quad \begin{cases} (\mathbf{T}^2 + \mathbf{n}\mathbf{T} + \text{id})(W)(u) = 0 \\ (\mathbf{T}_1 - \text{id})(W)(u) = 0 \end{cases} \quad (2-4)$$

2.4 Special 1-cut solutions

$\mathbf{T}$ is a linear operator on the space of meromorphic functions in the u -plane, over the field $\mathbf{k}$ of 1 and τ -translation invariant functions (the general form in the x variable of these bi-periodic functions is described Fig. 7). With the notation $\mathbf{n} = -2\cos(\pi\mu)$, the space of solutions of Eqn. 2-4 is the intersection of $\text{Ker}(\mathbf{T}_1 - \text{id})$ and:

$$\text{Ker}(\mathbf{T}^2 + \mathbf{n}\mathbf{T} + \text{id}) = \text{Ker}(\mathbf{T} - e^{i\pi\mu}\text{id}) \oplus \text{Ker}(\mathbf{T} - e^{-i\pi\mu}\text{id})$$

Thus, it has dimension two. Let us pick up [14, 18] in $\text{Ker}(\mathbf{T}_1 - \text{id}) \cap \text{Ker}(\mathbf{T} - e^{i\pi\mu}\text{id})$ a special function D_μ (i.e satisfying $D_\mu(u+1) = D_\mu(u)$ and $D_\mu(u+\tau) = e^{-i\pi\mu}D_\mu(u)$), having the asymptotic properties:

- $D_\mu(u(x)) \sim 1/x$ when $x \rightarrow \infty$ in the physical sheet.

- $D_\mu(u(x)) \propto 1/\sigma(x)$ when $x \rightarrow a, b, -a, -b$.

This determines D_μ uniquely, and we can actually construct it in terms of theta functions of modulus τ (see Appendix B for a reminder on theta functions):

$$D_\mu(u) = \frac{x(u)}{\sigma(x(u))} \frac{\vartheta_1\left(u - \frac{\tau}{2} + \frac{\mu}{2} \middle| \tau\right)}{\vartheta_1\left(u - \frac{\tau}{2} \middle| \tau\right)} \frac{\vartheta_1\left(-\frac{1}{2} \middle| \tau\right)}{\vartheta_1\left(-\frac{1}{2} + \frac{\mu}{2} \middle| \tau\right)} \quad (2-5)$$

By definition, it has two simple poles (mod $\mathbb{Z} \oplus \tau\mathbb{Z}$) at 0 and $\frac{1}{2}$, and a simple zero at $u(\infty) = -\frac{1}{2} + \frac{\tau}{2}$. It also admits a second simple zero at $u(e_\mu) = \frac{\tau-\mu}{2}$. This special function is our elementary brick to define a suitable basis of 1-cut solutions of Eqn. 2-4. Let us notice before that $u \mapsto D_\mu(-u)$ generates $\text{Ker}(\mathbf{T}_1 - \text{id}) \cap \text{Ker}(\mathbf{T} - e^{-i\pi\mu}\text{id})$.

To describe the space of 1-cut solutions of Eqn. 2-4, we rather work on the subfield $\mathbf{k}'$ of $\mathbf{k}$, consisting of u -even functions belonging to $\mathbf{k}$ (in the x variable, $\mathbf{k}'$ consists of rational functions of x^2). Then, this space of 1-cut solutions has again dimension two on $\mathbf{k}'$. We now describe a basis which we have found convenient for our purposes:

Theorem 2.1 *There exists a unique couple of meromorphic functions $(f_\mu, \widehat{f}_\mu)$, 1-cut solutions of Eqn. 2-4, such that:*

- $f_\mu(u(x))$ is holomorphic for $x \in \mathbb{C} \setminus [a, b]$ in the physical sheet (it has only one cut).
- $f_\mu(u(x)) \propto 1/\sigma(x)$ when $u \rightarrow u(a) = \tau$, or $u(b) = \frac{1}{2} + \tau$.
- $f_\mu(u(x)) \sim 1/x$ when $x \rightarrow \infty$ in the physical sheet.
- $\widehat{f}_\mu(u(x))$ is holomorphic for $x \in \mathbb{C} \setminus ([a, b] \cup \{0\})$ in the physical sheet (it has only one cut).
- $\widehat{f}_\mu(u)$ is finite when $u \rightarrow u(a)$ or $u(b)$.
- $\widehat{f}_\mu(u(x)) \propto 1/x$ when $x \rightarrow 0$ (in the two sheets).
- $\widehat{f}_\mu(u(x)) \sim 1$ when $x \rightarrow \infty$ in the physical sheet.

2.5 Bilinear form

The properties of these functions are listed in Appendix D. They were constructed such that $f_\mu \perp \widehat{f}_\mu = 0$ for the bilinear form (on the field $\mathbf{k}$) which appear in the loop equations (Thm. 1.2):

$$(g \perp h)(x) \equiv g(x)h(x) + g(-x)h(-x) + \frac{n}{2} (g(x)h(-x) + g(-x)h(x))$$

Equivalently in the u variable:

$$(g \perp h)(u) \equiv g(u)h(u) + g(u - \tau)h(u - \tau) + \frac{n}{2} (g(u)h(u - \tau) + g(u - \tau)h(u)) \quad (2-6)$$

Lemma 2.1

- If f and g are solutions of the homogeneous linear equation 2-2, then $(f \perp g)(u)$ is 1- and 2τ -translation invariant.
- Besides, if f and g have only one cut, then $f \perp g$ is an even rational function of x .
- If f is a 1-cut solution of 2-2, g has one cut, and $f \perp g$ has no cut on $[a, b]$, then g is a 1-cut solution of 2-2 as well.

proof:

For the first point, we rather work with the u variable than with x , and use the dictionary of Section 7: recall for example that "having one cut" translates into "being u -even". If f and g are solution of the loop equation:

$$\begin{aligned} \mathbf{T}(f \perp g) &= \mathbf{T}f \cdot \mathbf{T}g + \mathbf{T}^2 f \cdot \mathbf{T}^2 g + \frac{n}{2} (\mathbf{T}^2 f \cdot \mathbf{T}g + \mathbf{T}f \cdot \mathbf{T}^2 g) \\ &= (f \perp g) \end{aligned}$$

If on the top of that f and g have only one cut, so does $f \perp g$ for it is obviously invariant under $u \mapsto \tau - u$. Hence the second point. Now, assume f is a 1-cut solution of Eqn. 2-2 and $f \perp g$ has no cut on $[a, b]$ (for example, $f \perp g$ is an even rational function of x). Then, for $x \in [a, b]$ and $\epsilon \rightarrow 0^+$:

$$\begin{aligned} 0 &= (f \perp g)(x + i\epsilon) - (f \perp g)(x - i\epsilon) \\ &= \left(f(x + i\epsilon) + \frac{n}{2} f(-x) \right) g(x + i\epsilon) - \left(f(x - i\epsilon) + \frac{n}{2} f(-x) \right) g(x - i\epsilon) \\ &\quad + \frac{n}{2} (f(x + i\epsilon) - f(x - i\epsilon)) g(-x) \\ &= (f(x + i\epsilon) - f(x - i\epsilon)) \left[g(x + i\epsilon) + g(x - i\epsilon) + \frac{n}{2} g(-x) \right] \end{aligned}$$

Since f has a discontinuity on $]a, b[$, g must satisfy the homogeneous linear equation.

□

Thus, the norms $R_\mu(x^2) = (f_\mu \perp f_\mu)(x)$ and $\widehat{R}_\mu(x^2) = (\widehat{f}_\mu \perp \widehat{f}_\mu)(x)$ are rational functions of x^2 . They can be identified [14, 18] using the analytical properties of $(f_\mu, \widehat{f}_\mu)$ (see Appendix D).

2.6 General 1-cut solutions

We arrive at the description found in [14, 18].

Theorem 2.2 *The 1-cut solutions of Eqn. 2-2 are of the form*

$$W(x) = A(x^2) \frac{f_\mu(x)}{R_\mu(x^2)} + B(x^2) \frac{\widehat{f}_\mu(x)}{\widehat{R}_\mu(x^2)}$$

where $A(x^2)$ and $B(x^2)$ are rational functions of x^2 .

As rational functions of x^2 , A and B can be determined from the required analytic properties (behavior at poles and zeroes) of the solution of Eqn. 2-2 we are looking for.

3 Unstable generating functions ($2 - 2g - k \geq 0$)

We review the results of [14, 18, 19], concerning $W_1^{(0)}$ and $W_2^{(0)}$. They can be proved in a straightforward way from Thm. 2.2. Though the solution for $W_2^{(0)}$ requires lengthy computations. We have developed an other approach to compute $W_2^{(0)}$ (Section 3.4) which is closer to the spirit of algebraic geometry, that is, finding a good description of the ring of meromorphic functions on the spectral curve.

3.1 $W_1^{(0)}$ (rooted disks)

$W_1^{(0)}(x)$ is a 1-cut solution of the complete linear equation with RHS $V'(x)$. It is easy to find a particular solution of this equation:

$$V'_s(x) = \frac{2V'(x) - nV'(-x)}{4 - n^2}$$

So, $\overline{W}_1^{(0)}(x) = W_1^{(0)}(x) - V'_s(x)$ is a 1-cut solution of Eqn 2-2. It must satisfy:

- $W_1^{(0)}$ has no pole in the physical sheet (i.e in $\mathbb{C} \setminus [a, b]$).
- $W_1^{(0)}(x) \sim t/x$ when $x \rightarrow \infty$ in the physical sheet.
- $W_1^{(0)}(x)$ is finite when $x = a_i$ (a or b), and $W_1^{(0)}(x) - W_1^{(0)}(a_i) \propto \sqrt{(x - a_i)}$ when $x \rightarrow a_i$.

Theorem 3.1 [14, 18]

$$W_1^{(0)}(x) = \frac{2V'(x) - nV'(-x)}{4 - n^2} + A(x^2) \frac{f_\mu(x)}{R_\mu(x^2)} + B(x^2) \frac{\widehat{f}_\mu(x)}{\widehat{R}_\mu(x^2)} \quad (3-1)$$

where $A(x^2)$ and $B(x^2)$ are even polynomials:

$$\begin{aligned} A(x^2) &= -\frac{1}{2} \left(V'(x) f_\mu(x) + V'(-x) f_\mu(-x) \right)_+ \\ B(x^2) &= -\frac{1}{2} \left(V'(x) \widehat{f}_\mu(x) + V'(-x) \widehat{f}_\mu(-x) \right) \end{aligned}$$

a and b must be solution to the compatibility equations:

$$\begin{aligned} \operatorname{Res}_{x \rightarrow \infty} dx \left(x V'(x) f_\mu(x) \right) &= (-2 + \mathfrak{n}) t \\ \operatorname{Res}_{x \rightarrow \infty} dx \left(x V'(x) \widehat{f}_\mu(x) \right) &= (2 + \mathfrak{n}) \operatorname{Res}_{x \rightarrow \infty} dx \left(x W_1^{(0)}(x) \right) \end{aligned}$$

Generically, they admit a unique solution $\{a(t), b(t)\}$ which is a power series in $\sqrt{t}$ for small t (this $\sqrt{t}$ behavior is required by the 1-cut lemma).

Though $R_\mu(x)$ and $\widehat{R}_\mu(x)$ have a zero when $x^2 = e_\mu^2$, $W_1^{(0)}$ must be regular at $x = \pm e_\mu$, and this is expressed by the second compatibility equation.

The polynomial parts can be rewritten by taking residues at ∞ . By moving the contour, we can represent the solution as a contour integral around $[a, b]$ for the solution.

Theorem 3.2 [14, 18]

$$W_1^{(0)}(x_0) = \frac{1}{2i\pi} \oint_{[a,b]} dx \frac{x V'(x)}{x_0^2 - x^2} \left(\frac{f_\mu(x_0)}{R_\mu(x_0^2)} f_\mu(x) + \frac{\widehat{f}_\mu(x_0)}{\widehat{R}_\mu(x_0^2)} \widehat{f}_\mu(x) \right)$$

This formula is the analog for $\mathfrak{n} \neq 0$ of Tricomi's solution [39].

3.2 $y(x)$ (the spectral curve)

The object which plays the role of a spectral curve is the discontinuity of the resolvent $W_1^{(0)}(x)$. We have:

$$y(x) \equiv \lim_{\epsilon \rightarrow 0} \left(W_1^{(0)}(x + i\epsilon) - W_1^{(0)}(x - i\epsilon) \right) = 2W_1^{(0)}(x) + \mathfrak{n}W_1^{(0)}(-x) - V'(x)$$

A short computation with help of Appendix D gives:

$$y(x) = A(x^2) y_\mu(x) + B(x^2) \widehat{y}_\mu(x)$$

where the two basic blocks are given by:

$$y_\mu(x) = \frac{x\sigma(x)}{x^2 - e_\mu^2} \widehat{f}_\mu(-x), \quad \widehat{y}_\mu(x) = -\frac{x\sigma(x)}{x^2 - e_\mu^2} f_\mu(-x)$$

3.3 $W_2^{(0)}$ (rooted cylinders), first method

$W_2^{(0)}(x_0, x)$ is obtained by application of the loop insertion operator $\partial/\partial V(x)$ to $W_1^{(0)}(x_0)$. Accordingly, it is a 1-cut solution (in both x_0 and x) of the complete linear equation with RHS $-1/(x_0 - x)^2$. Again, we can decompose into a particular solution, and a solution $\overline{W}_2^{(0)}$ of Eqn. 2-2:

$$W_2^{(0)}(x_0, x) = \frac{-2}{(x + x_0)^2} + \frac{\mathbf{n}}{(x + x_0)^2} + \overline{W}_2^{(0)}(x_0, x)$$

We must find the solution characterized by:

- (i) $W_2^{(0)}(x_0, x)$ is symmetric in x_0 and x .
- (ii) $W_2^{(0)}(x_0, x)$ has no pole when x_0 and x are in the physical sheet.
- (iii) $W_2^{(0)}(x_0, x) \propto 1/x_0^2$ when $x_0 \rightarrow \infty$ in the physical sheet (x fixed).
- (iv) $W_2^{(0)}(x_0, x) \propto 1/\sqrt{(x_0 - a)(x_0 - b)}$ when $x_0 \rightarrow a, b$ (x fixed).

Indeed: (i) expresses the invariance of the generating function of cylinder maps by exchange of the marked faces ; (ii) is implied by the 1-cut lemma ; (iii) comes from the definition of $W_2^{(0)}$ as a connected correlation function ; (iv) follows from application of $\partial/\partial V(x)$ to $W_1^{(0)}(x_0)$, given that a and b are V dependent. Actually, it is easier to determine first the primitive of $\overline{W}_2^{(0)}$

$$H(x_0, x) = \int_{\infty}^x d\xi \overline{W}_2^{(0)}(x_0, \xi) \quad (3-2)$$

which satisfy assertions deduced from (i) – (iv). Although it is convenient to use the 1-cut basis $(f_{\mu}(x), \widehat{f}_{\mu}(x))$ to derive the result, it can be expressed afterwards in another basis. Let us introduce:

$$I(x) = \frac{x\sigma(x) + e_{\mu}\sigma(e_{\mu})}{x^2 - e_{\mu}^2} \quad (3-3)$$

which is related to the derivative of D_{μ} (see Appendix D).

Theorem 3.3 [14, 18] *Primitive of $W_2^{(0)}$.*

$$\begin{aligned} H(x_0, x) &= \frac{1}{4 - \mathbf{n}^2} (e^{-i\pi\mu} H_+(x_0, x) + H_+(-x_0, x) - H_+(x_0, -x) - e^{i\pi\mu} H_+(-x_0, -x)) \\ H_+(x_0, x) &\equiv D_{\mu}(x_0) \sigma(x) D_{\mu}(x) \frac{I(x) - I(x_0)}{x^2 - x_0^2} \end{aligned}$$

Corollary 3.1

$$\begin{aligned}
W_2^{(0)}(x_0, x) &= \frac{1}{4 - \mathbf{n}^2} \left(e^{-i\pi\mu} W_{2|+}^{(0)}(x_0, x) + W_{2|+}^{(0)}(-x_0, x) \right. \\
&\quad \left. + W_{2|+}^{(0)}(x_0, -x) + e^{i\pi\mu} W_{2|+}^{(0)}(-x_0, -x) \right) \\
W_{2|+}^{(0)}(x_0, x) &\equiv D_\mu(x_0) D_\mu(x) \left[1 - \left(\alpha_1 + \frac{x\sigma(x) - x_0\sigma(x_0)}{x^2 - x_0^2} \right) \frac{I(x) - I(x_0)}{x^2 - x_0^2} \right] - \frac{1}{(x + x_0)^2}
\end{aligned}$$

We also have:

$$\begin{aligned}
W_2^{(0)}(x, x) &= \frac{1}{4 - \mathbf{n}^2} \left(\frac{\mathbf{n}}{4x^2} - \mathbf{n}(\partial_2 H)(x, x) + 2W_{2|+}^{(0)}(x, -x) \right) \\
W_{2|+}^{(0)}(x, -x) &= \frac{1}{2} \frac{e_\mu^2 - \alpha_1^2}{\sigma^2(x)} + \frac{x^2 (b^2 - a^2)^2}{4 \sigma^4(x)}
\end{aligned}$$

3.4 $W_2^{(0)}$, infinite series representation

3.4.1 Decomposition and characterization

We present here an alternative construction of $W_2^{(0)}$, which takes advantage of the change of variable $x \rightarrow u(x)$. The natural object to look at is the differential form $W_2^{(0)}(x_0, x) dx_0 dx$. Let us define $s(u) = \frac{dx}{du} = \frac{ib}{2K'} \sigma(x(u))$. Its properties are:

$$s(u) = s(u + 1), \quad s(-u) = -s(u), \quad s(\tau - u) = s(u)$$

We want to determine:

$$\bar{\omega}_2^{(0)}(u, u_0) = s(u) s(u_0) \bar{W}_2^{(0)}(u, u_0)$$

We shall consider the function $\mathcal{B}$ defined by:

$$\mathcal{B}(u_0, u) \stackrel{\text{def}}{=} -e^{-i\pi\mu} \bar{\omega}_2^{(0)}(u, u_0) + \bar{\omega}_2^{(0)}(\tau - u, u_0) + \bar{\omega}_2^{(0)}(u, \tau - u_0) - e^{i\pi\mu} \bar{\omega}_2^{(0)}(\tau - u, \tau - u_0)$$

Conversely, $\bar{\omega}_2^{(0)}$ can be decomposed in terms of $\mathcal{B}$:

$$\bar{\omega}_2(u, u_0) = \frac{1}{4 - \mathbf{n}^2} (e^{-i\pi\mu} \mathcal{B}(u, u_0) + \mathcal{B}(\tau - u, u_0) + \mathcal{B}(u, \tau - u_0) + e^{i\pi\mu} \mathcal{B}(\tau - u, \tau - u_0)) \quad (3-4)$$

$\mathcal{B}$ has the following properties:

- $\mathcal{B}$ is a meromorphic function of $u, u_0 \in \mathbb{C}$.
- $\mathcal{B}$ has for only pole $u = (\tau - u_0) \bmod \mathbb{Z} \oplus \tau\mathbb{Z}$, and:

$$\mathcal{B}(u) = \frac{1}{(u + u_0 - \tau)^2} + O(1) \quad \text{when } u \rightarrow \tau - u_0.$$

- $\mathcal{B}(u, u_0) = \mathcal{B}(u_0, u)$.
- $\mathcal{B}(u, u_0) = \mathcal{B}(u + 1, u_0)$ and $\mathcal{B}(u + \tau, u_0) = e^{i\pi(1-\mu)} \mathcal{B}(u, u_0)$.

3.4.2 Construction of a solution

We can construct such a function explicitly by a deformation of the Weierstraß function $\wp$ (see Appendix C for a reminder of $\wp$). We assume $e^{i\pi\mu} \neq -1$ (i.e. $\mu \neq 2$), and define a function $\wp_\mu$:

$$\wp_\mu(w) = \sum_{m \in \mathbb{Z}} e^{i\pi(\mu-1)m} \frac{\pi^2}{\sin^2 \pi(w + m\tau)} \quad (3-5)$$

Notice that the series converges absolutely for $\tau \in i\mathbb{R}_+^*$. $\wp_\mu$ is 1-translation invariant and take a phase $e^{i\pi(1-\mu)}$ when $w \mapsto w + \tau$. It coincides with $\wp + c_0$ when $\mu = 1 \bmod 2\mathbb{Z}$ (i.e. $\mu = 2$), where c_0 is a constant depending on τ . It has for only pole $w = 0 \bmod (\mathbb{Z} \oplus \tau\mathbb{Z})$, which is a double pole without residues. As for D_μ , one can represent $\wp_\mu$ as a quotient of theta functions, and we find that it has two distinct zeroes :

$$w_1 \quad \text{and} \quad w_2 = -w_1 + \frac{1-\mu}{2}$$

$\wp_\mu$ enjoys properties generalizing those of $\wp$ and presented in Appendix C. Now, $\mathcal{B}(u, u_0) - \wp_\mu(u + u_0 - \tau)$ is holomorphic, 1-translation invariant and takes a phase $e^{i\pi(1-\mu)}$ when one of the arguments is shifted by τ . Since $\mu \in \mathbb{R}$, this difference is a entire bounded function⁶, so is constant, and this constant must vanish for $e^{i\pi\mu} \neq -1$. Therefore, $\mathcal{B}(u, u_0) = \wp_\mu(u + u_0 - \tau)$. Collecting the terms of Eqn. 3-4:

Theorem 3.4

$$\bar{w}_2^{(0)}(u, u_0) = \sum_{m \in \mathbb{Z}} \frac{2\pi^2 \cos(\pi(1-\mu)m)}{4 - \mu^2} \left(\frac{1}{\sin^2 \pi(u - u_0 + m\tau)} - \frac{1}{\sin^2 \pi(u + u_0 + m\tau)} \right)$$

Compared to Cor. 3.1, this expression is well suited if one wishes to integrate $W_2^{(0)}(x_0, x)$.

3.4.3 Comment

As usual in matrix models, we find that $\bar{w}_2^{(0)}(u_0, u)$ is a universal object. Universal means here that the dependence in V of $W_2^{(0)}$ arise only through a and b (it was already manifest in Cor. 3.1).

In the Coulomb gas picture (Eqn. 1-6), it characterizes the correlation of two eigenvalues in the large N limit. If we regard τ and $\mu \in \mathbb{R}/2\mathbb{Z}$ as independent parameters, Thm. 3.4 gives the Fourier series in μ of $\bar{W}_2^{(0)}$. Besides, it depends on $u(x) - u(x_0)$ and $u(x) - u(-x_0)$. This has a simple interpretation. Consider two eigenvalues located

⁶This argument is only valid when $\mu \in \mathbb{R}$, i.e requires at least $|\mu| \leq 2$.

at x_0 and x . In the $\mathcal{O}(\mathfrak{n})$ model, x_0 also feels the mirror charge located at $-x$. We found that the two point correlation function in the large N limit depends on the 'differences' between the position of the interacting sources. "Position" and "difference" have a natural meaning on the spectral curve, which is equipped with the addition law $(u_0, u) \mapsto u_0 + u$.

4 Stable generating functions ($2 - 2g - k < 0$)

We will see that the loop equations 1-10 imply that $W_k^{(g)}$ for $2g - 2 + k > 0$ satisfies the homogeneous linear equation 2-2. In principle, one could try to identify the even rational functions of x , $A_k^{(g)}(x^2, J)$ and $B_k^{(g)}(x^2, J)$ in a decomposition:

$$W_k^{(g)}(x, I) = A_k^{(g)}(x^2, I)f_\mu(x) + B_k^{(g)}(x^2, I)\widehat{f}_\mu(x)$$

This is basically the method of [1] to compute higher genus correlators, adapted to the $\mathcal{O}(\mathfrak{n})$ model in [18]. Inspired by [16], we shall take another route, which consists in finding a Cauchy residue formula for 1-cut solutions of Eqn. 2-2.

4.1 Cauchy residue formula

Also, we construct the appropriate Cauchy kernel $G(x_0, x)$ for our spectral curve:

$$\begin{aligned} G(x_0, x) &= \int_{-\infty}^x dx' \left(2\overline{W}_2^{(0)}(x_0, x') + \mathfrak{n}\overline{W}_2^{(0)}(x_0, -x') \right) \\ G(x_0, u) &= G(x_0, x(u)) \end{aligned} \quad (4-1)$$

It is possible to derive a infinite series representation of $s(u_0)s(u) G(x(u_0), x(u))$ from Thm. 3.4, but we will not use it. The essential properties of G are:

- $G(x_0, u)$ is a meromorphic function of $u \in \mathbb{C}$, with $u = u(x) \bmod (\mathbb{Z} \oplus 2\tau\mathbb{Z})$ as only pole.
- $G(x_0, x) = \frac{1}{x_0 - x} + O(1)$ when $x_0 \rightarrow x$.
- $G(x_0, -u) = -G(x_0, u)$.

Theorem 4.1 *Cauchy formula. Let W (in the x -plane) or W (in the u -plane) be a 1-cut solution of Eqn. 2-2. If W is holomorphic on $\mathbb{C} \setminus [a, b]$, and has no residue at $x = \infty$, we have:*

$$W(x_0) = \frac{1}{2} \operatorname{Res}_{u \rightarrow \tau, \tau + \frac{1}{2}} ds(u) G(x_0, u) W(u)$$

where $s(u) = \frac{dx}{du}$. One can add to G a constant $\gamma_i(x_0)$, depending on the branch point $u(a_i)$, $a_i \in \{a, b\}$, without changing the formula.

To prove this, we write the usual Cauchy formula and recast it as residues on the branch points a and b only.

$$\begin{aligned}
W(x_0) &= - \operatorname{Res}_{x \rightarrow x_0} dx G(x_0, x) W(x) \\
&= - \operatorname{Res}_{x \rightarrow x_0} dx \left(G(x_0, x) W(x) - \underbrace{G(x_0, -x) W(-x)}_{\text{no pole at } x_0} \right) \\
&= -\frac{1}{2} \operatorname{Res}_{x \rightarrow x_0, -x_0} dx (G(x_0, x) W(x) - G(x_0, -x) W(-x)) \quad \text{by parity}
\end{aligned}$$

We now switch to u variable: let $u_0 = u(x_0)$.

$$\begin{aligned}
W(x_0) &= -\frac{1}{4} \operatorname{Res}_{\substack{u \rightarrow u_0, -u_0 \\ \tau - u_0, -(\tau - u_0)}} du s(u) (G(x_0, u) W(u) - G(x_0, u - \tau) W(u - \tau)) \\
&= -\frac{1}{4} \operatorname{Res}_{\substack{u \rightarrow u_0, -u_0 \\ \tau - u_0, -(\tau - u_0)}} du (s(u) G(x_0, u) W(u) + s(u - \tau) G(x_0, u - \tau) W(u - \tau)) \\
&= -\frac{1}{4} \operatorname{Res}_{\substack{u \rightarrow u_0, -u_0 \\ \tau - u_0, -(\tau - u_0)}} du (s H(x_0, \cdot) \perp W)(u)
\end{aligned}$$

where H is defined in Eqn. 3-2. One knows that $s(u)H(x_0, u)$ (up to a constant in u , irrelevant in the residue) satisfies the linear equation in u . So, Lemma 2.1 tells us that $(sH(x_0, \cdot) \perp W)(u)$ is 1- and 2τ -translation invariant, i.e is elliptic. Since the sum of residues of an elliptic function vanishes, we have:

$$W(x_0) = \frac{1}{4} \operatorname{Res}_{\substack{u \rightarrow 0, \frac{1}{2} \\ \tau, \tau + \frac{1}{2}}} du (s H(x_0, \cdot) \perp W)(u)$$

Eventually, the assumption of u -parity/1-cut property for W allows us to rewrite:

$$W(x_0) = \frac{1}{2} \operatorname{Res}_{u \rightarrow \tau, \tau + \frac{1}{2}} du s(u) G(x_0, u) W(u)$$

4.2 $W_k^{(g)}$ (all topologies)

4.2.1 Recursive residue formula for $2g - 2 + k > 0$

In the matrix model, $W_k^{(g)}(x_1, \dots, x_k)$ represents correlation of densities of eigenvalues, hence have to be integrated on some interval of $[a, b]$ to give physical results. It is natural to look at the differential forms:

$$dx_1 \cdots dx_k W_k^{(g)}(x_1, \dots, x_k)$$

These forms can be pushed backwards under $u \mapsto x(u)$ to define differential forms $\omega_k^{(g)}$ on the u -domain $]-\frac{1}{2}, \frac{1}{2}[\times] - \tau, \tau[$.

$$\begin{aligned}\omega_k^{(g)}(u_1, \dots, u_k) &= du_1 s(u_1) \cdots du_k s(u_k) W(u_1, \dots, u_k) \\ &= dx(u_1) \cdots dx(u_k) W(x(u_1), \dots, x(u_k))\end{aligned}$$

and we recall that $s(u) = \frac{dx}{du}$. The main result of this article is:

Theorem 4.2 *For $2g - 2 + k > 0$, $W_k^{(g)}$ is a 1-cut solution of the homogeneous linear equation in each variable:*

$$W_k^{(g)}(x + i\epsilon, I) + nW_k^{(g)}(-x, I) + W_k^{(g)}(x - i\epsilon, I) \underset{\epsilon \rightarrow 0}{=} 0$$

Thus, $\omega_k^{(g)}(u_1, \dots, u_k)$ can be extended for $u_1, \dots, u_k \in \mathbb{C}$, as a meromorphic form in each u_i , with poles only at points where $x(u_i) = a$ or b . We have the recursive formula:

$$\omega_k(u_0, I) = \operatorname{Res}_{u \rightarrow \tau, \tau + \frac{1}{2}} du \mathcal{K}(u_0, u) \left[\omega_{k+1}^{(g-1)}(u, \bar{u}, I) + \sum'_{J \subseteq I, 0 \leq h \leq g} \bar{\omega}_{|J|+1}^{(h)}(u, J) \bar{\omega}_{k-|J|}^{(g-h)}(\bar{u}, {}^c J) \right]$$

(4-2)

- $\sum'_{(J,h)}$ is a sum over $J \subseteq I, 0 \leq h \leq g$, excluding $(J, h) = (\emptyset, 0)$ and (I, g) .
- $\bar{u} = 2\tau - 1 - u$ is such that $x(u) = x(\bar{u})$.
- The recursion kernel $\mathcal{K}(u_0, u)$ is a differential form in u , and the inverse of a differential form in u , given by:

$$\mathcal{K}(u_0, u) = -\frac{1}{2} \frac{\int_{\bar{u}}^u \bar{\omega}_2^{(0)}(u_0, u')}{(ydx)(u)} \quad (4-3)$$

- $\omega_{k'}^{(g')}(u, \bar{u}, I)$ differs from $\omega_{k'}^{(g')}$ only for $(k', g') = (2, 0)$, and:

$$\omega_2^{(0)}(u, \bar{u}) = - \left(\omega_2^{(0)}(u, u) + n\omega_2^{(0)}(u, \tau - u) + \omega_2^{(0)}(\tau - u, \tau - u) \right) \quad (4-4)$$

- When $(k, g) \neq (1, 1)$, the same formula is also true if one replaces $\omega_{k+1}^{(g-1)}(u, \bar{u}, I)$ by $\omega_{k+1}^{(g-1)}(u, u, I)$. However, one cannot in general replace $\bar{u}$ by u in the terms of $\sum'_{J,h}$.

4.2.2 Comment

We have found that the generating functions for genus g maps, carrying closed loops, with k boundaries, are given by the same recursive structure than the generating function for maps without decorations. Such a structure is called "topological recursion" [20]. It is a general fact that the counting of stable⁷ maps is more uniform than the counting of the unstable ones (rooted disks and rooted cylinders). This result provides a recursive algorithm to compute these generating functions. Minus the Euler characteristics of maps $|\chi| = 2g - 2 + k$ decreases by 1 at each step, and $W_k^{(g)}$ is reached with a stack $2g - 2 + k$ residues.

This result provides a first example where a topological recursion holds for a model which admits a non-algebraic plane curve as spectral curve:

$$\mathcal{L}_n = (\mathcal{T} = \mathbb{C}/(\mathbb{Z} \oplus \tau\mathbb{Z}), x, y)$$

$n = -2 \cos(\pi\mu)$ is the deformation parameter. The set of branch points is $\{0, \frac{1}{2}\} \subseteq \mathcal{T}$. But $y(u)$, $\mathcal{K}(u_0, u)$, $\dots$ are multivalued function on $\mathcal{T}$, basically constructed with the function D_μ which takes a phase $e^{i\pi\mu}$ under τ -translation. We shall give a univocal meaning to the notions of "function" or "differential form on the spectral curve" in Section 5.3, where we need it for computations. Though, we keep their polysemy in the next paragraph.

The initial data to run the algorithm of the topological recursion is not only a plane curve $(\mathcal{C}, x, y)$, but also of a Bergman kernel⁸ which is closely related to $\bar{\omega}_2^{(0)}$. For example, in the 1-matrix model with two cuts, the plane curve associated to the model is a torus, and the Weierstrass function $\wp(u - u_0) + \text{cte}$ are the possible Bergman kernel of the curve. Here, the notion of Bergman kernel is slightly deformed, and one may consider the Weierstrass $\wp_\mu$ function as the appropriate Bergman kernel. The recursion kernel should always be given by:

$$\mathcal{K}(u, u_0) = -\frac{1}{2} \frac{\int_{\bar{u}}^u \bar{\omega}_2^{(0)}(\cdot, u_0)}{(ydx)(u)}$$

The introduction of $\bar{u}$ (especially in the bracket term of Eqn. 4-2) is essential in the formulation of the topological recursion. If one try to replace $\bar{u}$ by u , the expression becomes wrong outside of the 1 hermitian matrix model. The reason of being of this $\bar{u}$ begins to be understood in the geometrical interpretation of the topological recursion [22].

⁷A Riemann surface with k marked points and genus g is called stable if it has only a finite number of automorphisms. This happens iff $2g - 2 + k > 0$. We carry this notion to discrete maps and correlation function.

⁸Recall that a Bergman kernel $du_0 du \mathcal{B}(u_0, u)$ is a meromorphic form on the spectral curve with a pole only when $x(u) = x(u_0)$, or order 2 and without residues.

4.2.3 Proof

$W_k^{(g)}$ satisfies the loop equation:

$$2 \left(W_k^{(g)}(\cdot, I) \perp \overline{W}_1^{(0)} \right) (x) = -E_k^{(g)}(x, I) - Q_k^{(g)}(x, I)$$

where

$$\begin{aligned} E_k^{(g)}(x, I) &= W_{k+1}^{(g-1)}(x, x, I) + W_{k+1}^{(g-1)}(-x, -x, I) + n W_{k+1}^{(g-1)}(x, -x, I) \\ &\quad + \sum_{J \subseteq I, 0 \leq h \leq g} (W_{|J|+1}^{(h)}(\cdot, J) \perp W_{k-|J|}^{(g-h)}(\cdot, {}^c J))(x) \\ Q_k^{(g)}(x, I) &= P_k^{(g)}(x, I) + P_k^{(g)}(-x, I) + \sum_{x_i \in I} \frac{\partial}{\partial x_i} \left[W_{k-1}^{(g)}(I) \left(\frac{1}{x - x_i} - \frac{1}{x + x_i} \right) \right] \end{aligned}$$

It is obvious that:

- $Q_k^{(g)}(\cdot, I)$ is finite when $x = a$ or b .
- $Q_k^{(g)}$ has no cut on $[a, b]$.

One can check easily as well (using symmetry in all variables of generating functions) that the combination of $W_{k+1}^{(g-1)}(\cdot, \cdot, I)$ entering in $E_k^{(g)}(x_0, I)$ has no cut on $[a, b]$. Since $Q_k^{(g)}$ has no cut on $[a, b]$ as well, an easy recursion on $2g - 2 + k$ and the second point in Lemma 2.1 show that $E_k^{(g)}(\cdot, I)$ has no cut on $[a, b]$. Hence, $(W_k^{(g)}(\cdot, I) \perp \overline{W}_1^{(0)})$ has no cut on $[a, b]$. Furthermore, according to the 1-cut lemma (Lemma 1.1), $W_k^{(g)}(\cdot, I)$ has only one cut. Now, the assumptions in the third point of Lemma 2.1 are satisfied, with $f = W_k^{(g)}(\cdot, I)$ and $g = \overline{W}_1^{(0)}$. So, $W_k^{(g)}(\cdot, I)$ must be a 1-cut solution of the homogeneous linear equation.

Now, we apply the Cauchy formula (Thm. 4.1), with arbitrary constants $\gamma_i(x_0)$:

$$\begin{aligned} &W_k^{(g)}(x_0, I) \\ &= \frac{1}{2} \operatorname{Res}_{u \rightarrow \tau, \tau + \frac{1}{2}} du s(u) (G(x_0, u) + \gamma_i(x_0)) W_k^{(g)}(u, I) \\ &= \frac{1}{2} \operatorname{Res}_{u \rightarrow \tau, \tau + \frac{1}{2}} du s(u) \frac{G(x_0, u) + \gamma_i(x_0)}{y(u)} \left(W_k^{(g)}(u, I) y(u) + \underbrace{W_k^{(g)}(u - \tau, I) y(u - \tau)}_{\text{no pole at } \tau, \tau + \frac{1}{2}} \right) \\ &= -\frac{1}{2} \operatorname{Res}_{u \rightarrow \tau, \tau + \frac{1}{2}} du s(u) \frac{G(x_0, u) + \gamma_i(x_0)}{y(u)} \left(E_k^{(g)}(u, I) + \underbrace{Q_k^{(g)}(u, I)}_{\text{is finite at } \tau, \tau + \frac{1}{2}} \right) \\ &= \operatorname{Res}_{u \rightarrow \tau, \tau + \frac{1}{2}} du \left[-\frac{1}{2} \frac{G(x_0, u) + \gamma_i(x_0)}{s(u) y(u)} \right] E_k^{(g)}(u, I) (s(u))^2 \end{aligned}$$

Besides, we have:

$$\begin{aligned}
G(x_0, u) &= \int_{u(\infty)}^u du' s(u') \left(2\overline{W}_2^{(0)}(x_0, u') + \mathbf{n}\overline{W}_2^{(0)}(x_0, \tau - u') \right) \\
&= \int_{u(\infty)}^u du' s(u') \left(\overline{W}_2^{(0)}(x_0, u') - \overline{W}_2^{(0)}(x_0, u' - 2\tau) \right) \\
&= \left(\int_{u(\infty)}^u du' s(u') \overline{W}_2^{(0)}(x_0, u') \right) - \left(\int_{2\tau - u(\infty)}^{2\tau - u} du' s(u') \overline{W}_2^{(0)}(x_0, u') \right) \\
&= \left(\int_{2\tau - u}^u du' s(u') \overline{W}_2^{(0)}(x_0, u') \right) + \left(\int_{u(\infty)}^{2\tau - u(\infty)} du' s(u') \overline{W}_2^{(0)}(x_0, u') \right) \\
\text{or} &= \left(\int_{2\tau + 1 - u}^u du' s(u') \overline{W}_2^{(0)}(x_0, u') \right) + \left(\int_{u(\infty)}^{2\tau - u(\infty)} du' s(u') \overline{W}_2^{(0)}(x_0, u') \right) \\
&\quad - \left(\int_0^1 du' s(u') \overline{W}_2^{(0)}(x_0, 2\tau - u') \right)
\end{aligned}$$

We used the 1-translation invariance for the last line. We see that, in a neighborhood of $u(a_i) = \tau$ (resp. $u(a_i) = \tau + \frac{1}{2}$):

$$G(x_0, u) = \int_{\bar{u}}^u du' s(u') \overline{W}_2^{(0)}(x_0, u') + \text{cte}_i(x_0)$$

and we can cancel this constant by the choice of $\gamma_i(x_0)$. Thus, the quantity in bracket (the recursion kernel) is indeed given by Eqn. 4-3 (written completely in variable u).

We assume now $(k, g) \neq (1, 1)$. Some terms in $E_k^{(g)}$ have $u - \tau$ as arguments, hence are regular at $u(a) = \tau$ and $u(b) = \tau + \frac{1}{2}$ and do not contribute to the residue. We make use of the homogeneous linear equation to write them apart. Since $(k+1, g-1) \neq (2, 0)$, $W_{k+1}^{(g-1)}(u, \bar{u}, I)$ exists, and:

$$\begin{aligned}
E_k^{(g)}(u, I) &= -W_{k+1}^{(g-1)}(u, \bar{u}, I) - \sum_{J \subseteq I, 0 \leq h \leq g} \overline{W}_{|J|+1}^{(h)}(u, J) \overline{W}_{k-|J|}^{(g-h)}(\bar{u}, {}^c J) \\
&\quad + \sum_{J \subseteq I, 0 \leq h \leq g} \overline{W}_{|J|+1}^{(h)}(u - \tau, J) \overline{W}_{k-|J|}^{(g-h)}(u - \tau, {}^c J) \\
&\quad + W_{k+1}^{(g-1)}(u - \tau, u - \tau, I)
\end{aligned}$$

Accordingly, we end up with:

$$W_k^{(g)}(u_0, I) = \text{Res}_{u \rightarrow \tau, \tau + \frac{1}{2}} du \mathcal{K}(u_0, u) \left[W_{k+1}^{(g)}(u, \bar{u}, I) + \sum_{J \subseteq I, 0 \leq h \leq g} \overline{W}_{|J|+1}^{(h)}(u, J) \overline{W}_{k-|J|}^{(g-h)}(\bar{u}, {}^c J) \right] s(u)(-s(u))$$

and we know that $s(\bar{u}) = -s(u)$. This can be rewritten with differential forms and we find Eqn. 4-2.

Eventually, we have to write separately the case $(k, g) = (1, 1)$, because the polar structure of $W_2^{(0)}$ is different. In $E_k^{(g)}(u, I)$, there is only one term, $-''W_2^{(0)}(u, \bar{u})''$, defined by

$$''W_2^{(0)}(u, \bar{u})'' \equiv - \left(W_2^{(0)}(u, u) + \mathbf{n}W_2^{(0)}(u, \tau - u) + W_2^{(0)}(\tau - u, \tau - u) \right)$$

If we set:

$$\begin{aligned} ''\omega_2^{(0)}(u, \bar{u})'' &\equiv -(\mathrm{d}x(u))^2 \left(W_2^{(0)}(u, u) + \mathbf{n}W_2^{(0)}(u, \tau - u) + W_2^{(0)}(\tau - u, \tau - u) \right) \\ &= - \left(\omega_2^{(0)}(u, u) + \mathbf{n}\omega_2^{(0)}(u, \tau - u) + \omega_2^{(0)}(\tau - u, \tau - u) \right), \end{aligned}$$

then Eqn. 4-2 is still correct.

4.3 Examples

Finding $W_3^{(0)}$ and $W_1^{(1)}$ involve only one residue computation. Let us call v_i the branch points $u(a) = \tau$ and $u(b) = \tau + \frac{1}{2}$. We shall use the following notation:

$$\begin{aligned} \bar{\omega}_2^{(0)}(u_0, v_i) &\equiv \left(\frac{\bar{\omega}_2^{(0)}(u_0, u)}{\mathrm{d}u} \right) \Big|_{u=v_i} \\ \text{and } (\partial_2^m \bar{\omega}_2^{(0)})(u_0, v_i) &\equiv \frac{\partial^m}{\partial u^m} \left(\frac{\bar{\omega}_2^{(0)}(u_0, u)}{\mathrm{d}u} \right) \Big|_{u=v_i} \quad \text{for } m \geq 1 \end{aligned}$$

I.e., we identify differential forms and families of functions indexed by an atlas of $\mathbb{C}$ and having the proper transformation under a change of local coordinate, and we read all differential forms in the local coordinate u .

4.3.1 Recursion kernel at the branchpoints

As a preliminary to the computations, we give the Laurent expansion of $\mathcal{K}(u, u_0)$ when $\delta = u - v_i \rightarrow 0$. In the local coordinate u :

$$\begin{aligned} \mathcal{K}(u_0, v_i + \delta) &= -\frac{\bar{\omega}_2^{(0)}(u_0, v_i)}{y'(v_i)s'(v_i)} \frac{1}{\delta} + \\ &\quad \left[\frac{1}{4} \frac{\bar{\omega}_2^{(0)}(u_0, v_i)}{y'(v_i)s'(v_i)} \left(\frac{y'''(v_i)}{y'(v_i)} + \frac{s'''(v_i)}{s'(v_i)} \right) - \frac{1}{6} \frac{\partial_2^2 \bar{\omega}_2^{(0)}(u_0, v_i)}{y'(v_i)s'(v_i)} \right] \delta + o(\delta) \end{aligned} \tag{4-5}$$

We notice $\mathcal{K}(u_0, v_i + \delta)$ is odd in δ since $\mathcal{K}$ is changed in $-\mathcal{K}$ when $u \leftrightarrow \bar{u}$. All the same, only the odd order derivative (with respect to u) of s and y do not vanish a priori at $u = v_i$.

4.3.2 $W_3^{(0)}$ (pairs of pants)

Theorem 4.3

$$\omega_3^{(0)}(u_0, u_1, u_2) = \sum_i \frac{-2}{y'(v_i)s'(v_i)} \bar{\omega}_2^{(0)}(u_0, v_i) \bar{\omega}_2^{(0)}(u_1, v_i) \bar{\omega}_2^{(0)}(u_2, v_i)$$

4.3.3 $W_1^{(1)}$ (rooted disk with one handle)

$W_1^{(1)}$ is the generating function for rooted toroidal maps. According to the topological recursion:

$$\omega_1^{(1)}(u_0) = \sum_i \operatorname{Res}_{u \rightarrow v_i} \mathcal{K}(u, u_0) {}^{\prime\prime}\omega_2^{(0)}(u, \bar{u}){}^{\prime\prime}$$

Using the Laurent expansion of $\mathcal{K}(u_0, u)$ when $u \rightarrow v_i$ (Eqn. 4-5), we find:

$$\begin{aligned} \omega_1^{(1)}(u_0) &= -\frac{\bar{\omega}_2^{(0)}(u_0, v_i)}{y'(v_i)s'(v_i)} \operatorname{Res}_{u \rightarrow v_i} \frac{{}^{\prime\prime}\omega_2^{(0)}(u, \bar{u}){}^{\prime\prime}}{u - v_i} \\ &\quad + \frac{1}{4} \frac{\bar{\omega}_2^{(0)}(u_0, v_i)}{y'(v_i)s'(v_i)} \left(\frac{y'''(v_i)}{y'(v_i)} + \frac{s'''(v_i)}{s'(v_i)} \right) \operatorname{Res}_{u \rightarrow v_i} (u - v_i) {}^{\prime\prime}\omega_2^{(0)}(u, \bar{u}){}^{\prime\prime} \\ &\quad - \frac{1}{6} \frac{\partial_2^2 \bar{\omega}_2^{(0)}(u_0, v_i)}{y'(v_i)s'(v_i)} \operatorname{Res}_{u \rightarrow v_i} (u - v_i) {}^{\prime\prime}\omega_2^{(0)}(u, \bar{u}){}^{\prime\prime} \end{aligned}$$

Besides, we can make use of the representation of $\bar{\omega}_2^{(0)}$ as an infinite series (Thm. 3.4) and compute explicitly:

$$\begin{aligned} \frac{{}^{\prime\prime}\omega_2^{(0)}(u, \bar{u}){}^{\prime\prime}}{(du)^2} &= -\frac{1}{(du)^2} \left(\omega_2^{(0)}(u, u) + n\omega_2^{(0)}(u, \tau - u) + \omega_2^{(0)}(\tau - u, \tau - u) \right) \\ &= \lim_{u \rightarrow u_0} \left(\frac{s(u)s(u_0)}{(x(u) - x(u_0))^2} - \sum_{m \in \mathbb{Z}} \frac{(2r_m - nr_{m+1})\pi^2}{\sin^2 \pi(u - u_0 + m\tau)} \right) \\ &= \frac{1}{6} (Sx)(u) - \frac{\pi^2}{3} + \sum_{m \geq 1} \frac{\pi^2 \cos \pi(1 - \mu)m}{\sin^2 \pi m\tau} \end{aligned}$$

We have introduced the schwartzian derivative of x with respect to u :

$$(Sx)(u) \equiv \frac{x'''(u)}{x'(u)} - \frac{3}{2} \left(\frac{x''(u)}{x'(u)} \right)^2$$

When $u \rightarrow v_i$, the first term is finite whereas the second one has a double pole:

$$(Sx)(u) = -\frac{3}{2} \frac{1}{(u - v_i)^2} + O((u - v_i)^2) \quad \text{when } u \rightarrow v_i$$

Hence, $\text{Res}_{u \rightarrow v_i} (u - v_i)'' \omega_2^{(0)}(u, \bar{u})'' = -\frac{1}{4}$. Eventually, let us introduce the so-called *connective projection*:

$$\begin{aligned} S_{B,x}(v_i) &\stackrel{\text{def}}{=} -6 \text{Res}_{u \rightarrow v_i} \frac{''\omega_2^{(0)}(u, \bar{u})''}{dx(u)} \\ &= \frac{1}{s'(v_i)} \left(2\pi^2 + \sum_{m \in \mathbb{Z} \setminus \{0\}} \frac{6q_m \pi^2}{\sin^2 \pi m \tau} \right) \end{aligned}$$

We have in terms of $S_{B,x}$:

$$\text{Res}_{u \rightarrow v_i} \frac{''\omega_2^{(0)}(u, \bar{u})''}{(u - v_i)} = -\frac{1}{6} S_{B,x}(v_i)$$

The final result is:

Theorem 4.4

$$\omega_1^{(1)}(u_0) = \sum_i \frac{1}{24} \frac{(\partial_2^2 \bar{\omega}_2^{(0)})(u_0, v_i)}{y'(v_i) s'(v_i)} + \frac{\bar{\omega}_2^{(0)}(u_0, v_i)}{y'(v_i)} \left(\frac{S_{B,x}(v_i)}{6} - \frac{1}{16} \frac{s'''(v_i)}{s'(v_i)^2} - \frac{1}{16} \frac{y'''(v_i)}{y'(v_i) s'(v_i)} \right)$$

This expression is, again, similar to the 1-matrix model. We have given the details of the computation to illustrate that the steps are the same as in the 1-matrix model, provided the definition of $''\omega_2^{(0)}(u, \bar{u})''$ is adapted to the $\mathcal{O}(\mathfrak{n})$ model according to Eqn. [4-4](#).

5 Properties of the $W_k^{(g)}$'s

In this section, we show that the properties found with the topological recursion in the 1-matrix model can be completely generalized for the deformed topological recursion relevant in the $\mathcal{O}(\mathfrak{n})$ model. The special geometry structure is present, the stable F_g 's ($g \geq 2$) are computed by the same integration formula, and (almost) the same formula exists for F_0 and F_1 .

Once the properties for unstable quantities are checked, many proofs can be done by induction like in the 1-hermitian matrix model. Those proofs only use the residue recursion formula Eqn. [4-2](#) and a few other simple properties which are satisfied here. We choose not to reproduce the derivation of these known results, for sake of brevity, and only stress the particularities in the $\mathcal{O}(\mathfrak{n})$ model. Also, we refer to [\[20\]](#) for the complete proofs.

5.1 Symmetry

Though it is not obvious, the residue formula Eqn. 4-2 yields symmetric $\omega_k^{(g)}$'s. This statement is based on the symmetry of $\omega_2^{(0)}$, and the loop equations Eqn. 1-10. The proof is similar to the case of the one hermitian matrix model.

5.2 Homogeneity

The partition function of the $\mathcal{O}(\mathbf{n})$ model (Eqn. 1-4), $Z(t, t_3, t_4, \dots)$, is invariant under

$$(t, t_0, t_1, t_2, t_3, \dots) \mapsto (\lambda t, \lambda t_0, \lambda t_1, \lambda t_2, \lambda t_3, \dots) \quad \lambda > 0$$

Therefore:

$$\forall k, g \geq 0, \quad \left(t \frac{\partial}{\partial t} + \sum_{j \geq 0} t_j \frac{\partial}{\partial t_j} \right) W_k^{(g)} = (2 - 2g - k) W_k^{(g)} \quad (5-1)$$

5.3 Special geometry

What we call special geometry is the data of a non degenerate pairing, which to a differential form $\widehat{\Omega}$ on the spectral curve, associates:

- a cycle $\Omega^* \subseteq \mathbb{C}$, i.e in the u -plane,
- a germ of holomorphic function Λ_Ω on Ω^* ,

with the following property, for all $k, g \geq 0$:

$$\begin{aligned} \delta_\epsilon \bar{\omega}_k^{(g)}(I) &= \int_{\Omega^*} du \Lambda_\Omega(u) \bar{\omega}_{k+1}^{(g)}(u, I) \\ \text{when } ydx &\rightarrow ydx + \epsilon \widehat{\Omega} \quad \epsilon \rightarrow 0 \end{aligned} \quad (5-2)$$

Though, for the $\mathcal{O}(\mathbf{n})$ model, we understand the notion of "differential form on the spectral curve" in a slightly deformed way. The manifold for the spectral curve can be considered as $\mathcal{T}$ or $\mathbb{C}$. If $\widehat{\Omega}$ is a differential meromorphic form on $\mathbb{C}$, we define:

$$\Omega = \frac{1}{4 - \mathbf{n}^2} \left(2\widehat{\Omega}(u) + \mathbf{n}\widehat{\Omega}(\tau - u) \right)$$

For us, $\widehat{\Omega}$ is a differential form on the spectral curve if Ω/du is a u -even solution of Eqn. 2-4 (that is, $\widehat{\Omega}$ has the same properties as ydx). A consequence of Eqn. 5-2 is that the pairing is given by integration of $\bar{\omega}_2^{(0)}$ against Ω^* :

$$\begin{aligned} \Omega(u_0) &= \int_{\Omega^*} du \Lambda_\Omega(u) \bar{\omega}_2^{(0)}(u_0, u) \\ \widehat{\Omega}(u_0) &= \int_{\Omega^*} du \Lambda_\Omega(u) \left(2\bar{\omega}_2^{(0)}(u_0, u) - \mathbf{n}\bar{\omega}_2^{(0)}(\tau - u_0, u) \right) \end{aligned}$$

In the next paragraphs, we compute the variations of the $\omega_k^{(g)}$'s with respect to the t_j 's (which is a simple task) and to t (which gives an interesting result). Roughly speaking, we hold x fixed, but the value $u(x)$ changes since a and b are determined by the consistency relations and depend on t and on the t_j 's. To avoid confusions, we note $\partial_t|_x$, $\partial_{t_j}|_x$ these variations. By definition:

$$\left(\frac{\partial}{\partial t_\bullet}\right)_x \omega_k^{(g)}(u_1, \dots, u_k) = dx(u_1) \cdots dx(u_k) \left(\frac{\partial}{\partial t_\bullet} W\right)(x(u_1), \dots, x(u_k))$$

We show that for each of these parameters, one can find a dual cycle such that special geometry holds. If we had considered $W_1^{(0)}(x)$ with several cuts in the x plane, one would also like to compute variations with respect to filling fractions and identify the corresponding dual cycle.

5.3.1 Variations of the t_j 's

Theorem 5.1

$$\forall k, g \geq 0, \forall j \geq 3 \quad \left(\frac{\partial}{\partial t_j}\right)_x \omega_k^{(g)}(I) = \text{Res}_{u \rightarrow u(\infty)} du \frac{x(u)^j}{j} \omega_{k+1}^{(g)}(u, I)$$

The cycle associated to t_j is $(\Omega_j^*, \Lambda_j) = (\frac{1}{2i\pi} \mathcal{C}_\infty, \frac{x(u)^j}{j})$, where $\mathcal{C}_\infty$ is a contour surrounding $u(\infty)$ and no other special point.

We start from the definition of the correlation function before the topological expansion:

$$\begin{aligned} \left(\frac{\partial}{\partial t_1} W_k\right)(I) &= -\left\langle \frac{N}{t} \frac{\text{Tr} M^j}{j} \prod_{x_i \in I} \text{Tr} \frac{1}{x_i - M} \right\rangle_C \\ &= \text{Res}_{x \rightarrow \infty} dx \left\langle \frac{N}{t} \frac{x^j}{j} \text{Tr} \frac{1}{x - M} \prod_{x_i \in I} \text{Tr} \frac{1}{x_i - M} \right\rangle_C \end{aligned}$$

After expansion in powers of $\frac{N}{t}$, and translation into the language of differential forms, we obtain Thm. 5.1.

5.3.2 Variation of t

We begin with the obvious remark that $\partial_t|_x \bar{\omega}_k^{(g)} = \partial_t|_x \omega_k^{(g)}$. Let us compute first the variations of the stable correlation forms $\omega_1^{(0)}$ and $\omega_2^{(0)}$.

Theorem 5.2

$$\left(\frac{\partial}{\partial t}\right)_x \bar{\omega}_1^{(0)}(u) = \int_{u(\infty)}^{\overline{u(\infty)}} \bar{\omega}_2^{(0)}(\cdot, u)$$

We have defined the point $\overline{u(\infty)} = 2\tau - 1 - u(\infty) = \frac{-1+3\tau}{2}$. The cycle associated to t is $(\Omega_t^*, \Lambda_t) = ([\frac{-1+\tau}{2}, \frac{-1+3\tau}{2}], 1)$. There is a corresponding form on the spectral curve: $\Omega_t(u) = dx(u) f_\mu(u)$.

Theorem 5.3

$$\begin{aligned} \left(\frac{\partial}{\partial t} \Big|_{\mathbf{x}} \bar{\omega}_2^{(0)} \right) (u_1, u_2) &= \int_{\Omega_t^*} \omega_3^{(0)}(\cdot, u_1, u_2) \\ &= \sum_i \text{Res}_{u \rightarrow v_i} du \left[\mathcal{K}(u_1, u) \bar{\omega}_2^{(0)}(u, u_2) \left(\frac{\partial}{\partial t} \Big|_{\mathbf{x}} \bar{\omega}_1^{(0)} \right) (u) \right] + (u_1 \leftrightarrow u_2) \end{aligned} \quad (5-3)$$

To prove the formula 5-2 for $2g - 2 + k > 0$, we need first a lemma for the variation of the recursive kernel $\mathcal{K}$ (Eqn. 4-3).

Lemma 5.1

$$\left[\left(\frac{\partial}{\partial t} \Big|_{\mathbf{x}} + \frac{\partial \ln y}{\partial t}(u) \right) \mathcal{K} \right] (u_0, u) = \sum_i \text{Res}_{u' \rightarrow v_i} du' \left[\mathcal{K}(u_0, u') \mathcal{K}(u', u) \left(\frac{\partial}{\partial t} \Big|_{\mathbf{x}} \bar{\omega}_1^{(0)} \right) (u') \right]$$

Then, the general result follows:

Theorem 5.4 For all $k, g \neq (0, 0)$:

$$\left(\frac{\partial}{\partial t} \Big|_{\mathbf{x}} \bar{\omega}_k^{(g)} \right) (I) = \int_{\Omega_t^*} \bar{\omega}_{k+1}^{(g)}(\cdot, I)$$

This will be completed in Section 5.5 with expressions for the derivatives of F_0 .

Proof of Thm. 5.2

We derive the properties of $\partial_t|_{\mathbf{x}} \omega_1^{(0)}$ from those of $W_1^{(0)}$ listed in Section 3.1. They coincide with those of f_μ listed in Section 2.4, and both of these functions are 1-cut solution of the homogeneous linear equation. Hence $\left(\frac{\partial}{\partial t} \Big|_{\mathbf{x}} W_1^{(0)} \right) (\mathbf{x}) = f_\mu(\mathbf{x})$. Now, we would like to represent $f_\mu(x_0)dx_0$ as an integral of $\bar{\omega}_2^{(0)}$ over some path in the u -plane. Having a look at $H(x_0, x)$ in Thm. 3.3, it is possible to write:

$$f_\mu(u_0)dx(u_0) = \text{cte} \times \int_{\frac{-1+\tau}{2}}^{u^*} \bar{\omega}_2^{(0)}(\cdot, u_0) \quad (5-4)$$

if we take $u^* \neq u(\infty)$ such that $f_\mu(u^*) = 0$. An accurate value is:

$$u^* = \overline{u(\infty)} = 2\tau - 1 - u(\infty) = \frac{-1+3\tau}{2}$$

Using the homogeneous linear equation, one computes:

$$f_\mu(u_0) \sim \frac{n-1}{x(u_0)} \quad \text{when } u_0 \rightarrow \overline{u(\infty)}$$

and find that $f_\mu(u_0)dx(u_0) = H(u_0, \overline{u(\infty)})dx(u_0)$.

Proof of Thm. 5.3-Lemma 5.1

We compute with help successively of Eqn. 5-1, Thm. 5.1, Thm. 4.3, Thm. 5.2, and comparison to Eqn. 4-5:

$$\begin{aligned}
\left(t \frac{\partial}{\partial t} \Big|_x \bar{\omega}_2^{(0)}\right)(u_0, u_1) &= - \sum_j \left(t_j \frac{\partial}{\partial t_j} \Big|_x \bar{\omega}_2^{(0)}\right)(u_0, u_1) \\
&= - \operatorname{Res}_{u \rightarrow u(\infty)} V(x(u)) \omega_3^{(0)}(u, u_0, u_1) \\
&= 2 \sum_i \frac{\bar{\omega}_2^{(0)}(v_i, u_0) \bar{\omega}_2^{(0)}(v_i, u_1)}{y'(v_i) s'(v_i)} \operatorname{Res}_{u \rightarrow u(\infty)} \left(V(x(u)) \bar{\omega}_2^{(0)}(v_i, u)\right) \\
&= 2 \sum_i \frac{\bar{\omega}_2^{(0)}(v_i, u_0) \bar{\omega}_2^{(0)}(v_i, u_1)}{y'(v_i) s'(v_i)} \left[- \left(t \frac{\partial}{\partial t} \Big|_x \bar{\omega}_1^{(0)}\right)(v_i) + \bar{\omega}_1^{(0)}(v_i) \right] \\
&= -2 \sum_i \frac{\bar{\omega}_2^{(0)}(v_i, u_0) \bar{\omega}_2^{(0)}(v_i, u_1)}{y'(v_i) s'(v_i)} \left(t \int_{\frac{-1+\tau}{2}}^{\frac{-1+3\tau}{2}} \bar{\omega}_2^{(0)}(v_i, u) \right) \\
&= t \int_{\frac{-1+\tau}{2}}^{\frac{-1+3\tau}{2}} \omega_3^{(0)}(u, u_0, u_1) \\
\text{or } &= 2 \sum_i \operatorname{Res}_{u \rightarrow v_i} \left[\mathcal{K}(u_0, u) \bar{\omega}_2^{(0)}(u, u_1) \left(t \frac{\partial}{\partial t} \Big|_x \bar{\omega}_1^{(0)}\right)(u) \right]
\end{aligned}$$

The last line is an application of the lemma above. By symmetry of $(\partial_t|_x \bar{\omega}_2^{(0)})(u_0, u_1)$, we can also distribute u_0 and u_1 in two ways, as in Thm. 5.3.

Then, it is easy to prove Thm. 5.1 from the expression of $\partial_t|_x \bar{\omega}_2^{(0)}$ in the last line.

Proof of Thm. 5.4

For $(k, g) \neq (0, 0)$, the proof is similar to the 1-matrix model. We shall consider the case of $\partial_t F_0$ later, in Section 5.5.

5.4 Integration formula

The inverse operation, consisting in recovering $\omega_k^{(g)}$ from $\omega_{k+1}^{(g)}$, can also be performed by a residue calculation at the branch points.

Theorem 5.5 *For all k, g such that $2g - 2 + k \geq 0$:*

$$\boxed{\sum_i \operatorname{Res}_{u \rightarrow v_i} \phi(u) \omega_{k+1}^{(g)}(u, I) = (2 - 2g - k) \omega_k^{(g)}(I)} \tag{5-5}$$

where $d\phi = ydx$. In particular, for $g \geq 2$:

$$F_g = \frac{1}{2 - 2g} \sum_i \operatorname{Res}_{u \rightarrow v_i} \phi(u) \omega_1^{(g)}(u) \tag{5-6}$$

proof:

Similar to the 1-matrix model. $\square$ Though, F_0 and F_1 cannot be found by this method.

5.5 F_0 (spherical maps)

5.5.1 Earlier results

In the matrix model, F_0 is minus the free energy of the model in the thermodynamic limit. In other words, it gives the leading order of $\ln Z$ modulo the remark about convergent matrix integrals (Section 1.8). The saddle point technique applied to the partition function Eqn. 1-4 gives a heuristic formula:

Proposition 5.1

$$F_0 = -\frac{t}{2} \int_a^b dx \varrho(x) "V(x)"$$

where $\varrho(x) = -\frac{1}{2i\pi t} y(x)$ is the density of eigenvalues (which has $[a, b]$ as support on the real line) in the thermodynamic limit. " $V(x)$ " is the primitive of $V'(x)$ which satisfies:

$$\forall x_0 \in [a, b], \quad \int_a^b dx \rho(x) \left(2 \ln |x - x_0| - \frac{n}{2} \ln |x + x_0| \right) = "V(x_0)"$$

(It differs from $V(x)$ only by a constant term).

In fact, it is easier to compute derivatives of F_0 with respect to t . The most explicit expressions obtained in [19] were:

Theorem 5.6

$$\frac{\partial^3 F_0}{\partial t^3} = \frac{(2 - n)}{a^2 - b^2} \left[\frac{a^2 - e_\mu^2}{a} \frac{\partial a}{\partial t} - \frac{b^2 - e_\mu^2}{b} \frac{\partial b}{\partial t} \right] \quad (5-7)$$

It can be integrated once:

$$\frac{\partial^2 F_0}{\partial t^2} = \left(1 - \frac{n}{2} \right) \ln \left[a^2 g \left(1 - \frac{a^2}{b^2} \right) \right] + \hat{C}_2$$

where $\hat{C}_2$ is a constant independent of t , and g a function defined by:

$$\frac{d \ln g}{dm} = \frac{1}{m(1 - m) \operatorname{cd}_{\sqrt{m}}^2(\mu K(\sqrt{m}))}$$

and $\operatorname{cd}_k(w) = \frac{\operatorname{dn}_k(w)}{\operatorname{cn}_k(w)}$.

5.5.2 New results

With special geometry, we can compute $\partial_t^k F_0$ by integrating $\omega_k^{(0)}$ on $(\Omega_t^*)^k$. This is true for $k \geq 3$, but some adaptations are required for $k = 2$ and $k = 1$ because those integrals diverge. For these cases, we only give the result, which is very similar to the 1-hermitian matrix model. We recall that $s(u) = x'(u)$.

Theorem 5.7

$$\begin{aligned}\frac{\partial^2 F_0}{\partial t^2} &= \lim_{\epsilon \rightarrow 0} \left((2 - \mathbf{n}) \ln x(u(\infty) + \epsilon) + \int_{u(\infty) + \epsilon}^{u(b)} du \left((s f_\mu)(u) + (s f_\mu)(\bar{u}) \right) \right) + C_2 \\ \frac{\partial F_0}{\partial t} &= \lim_{\epsilon \rightarrow 0} \left((2 - \mathbf{n}) t \ln x(u(\infty) + \epsilon) - V(x(u(\infty) + \epsilon)) + \int_{u(\infty) + \epsilon}^{u(b)} du y(u) \right) \quad (5-8)\end{aligned}$$

By homogeneity (Eqn. 5-1), it is straightforward to obtain F_0 from Eqn. 5-8

Theorem 5.8

$$F_0 = \frac{1}{2} \operatorname{Res}_{x \rightarrow \infty} dx V(x) W_1^{(0)}(x) + \frac{t}{2} \frac{\partial F_0}{\partial t}$$

With the infinite series representation, we can compute further. As for $W_2^{(0)}$, $\partial_t^2 F_0$ depends on the potential V only through the parameters a and b , or equivalently through τ and K introduced in Section 2.2.

Theorem 5.9

$$\frac{\partial^2 F_0}{\partial t^2} = (2 - \mathbf{n}) \ln \left(\frac{\pi b |\tau|}{2K} \right) + 2\sqrt{4 - \mathbf{n}^2} \sum_{m \geq 1} T_m(\mathbf{n}/2) \ln (\sin \pi(1 - \mu)m)$$

$T_m(\mathbf{n}/2) = \cos \pi(1 - \mu)m$ is the m -th Chebyshev polynomial.

($\partial_t^2 F_0$ counts spherical maps with v vertices with a weight $v(v - 1)t^{v-2}$. In a rough way, we can say that this weight is proportional to the area of the dual map). The situation is very different for $\partial_t F_0$ and F_0 , which depends fully on the potential V . We have obtained an explicit formula for F_0 for any polynomial V , involving an infinite series. We do not reproduce it here for lisibility, but it is available on request.

5.6 F_1 (toroidal maps)

We can use special geometry to compute $\partial_t F_1$.

$$\begin{aligned}
\frac{\partial F_1}{\partial t} &= \int_{\Omega_t^*} \omega_1^{(1)} \\
&= \sum_i \frac{1}{24} \frac{\int_{u_0 \in \Omega_t^*} (\partial_2^2 \bar{\omega}_2^{(0)})(u_0, v_i)}{y'(v_i) s'(v_i)} + \\
&\quad + \frac{\int_{u_0 \in \Omega_t^*} \bar{\omega}_2^{(0)}(u_0, v_i)}{y'(v_i)} \left(\frac{S_B(v_i)}{6} - \frac{1}{16} \frac{s'''(v_i)}{s'(v_i)^2} - \frac{1}{16} \frac{y'''(v_i)}{y'(v_i) s'(v_i)} \right) \\
&= \sum_i \frac{1}{24} \frac{(sf_\mu)''(v_i)}{y'(v_i) s'(v_i)} + \frac{(sf_\mu)(v_i)}{y'(v_i)} \left(\frac{S_B(v_i)}{6} - \frac{1}{16} \frac{s'''(v_i)}{s'(v_i)^2} - \frac{1}{16} \frac{y'''(v_i)}{y'(v_i) s'(v_i)} \right)
\end{aligned}$$

The integration of the right hand side with respect to t has been studied in the literature [17], and relies on the computation of $\partial_t|_x(y'(v_i))$ and the comparison with Eqn. 5-9. This equation is essentially the same as the one encountered in the one matrix model. So, we give directly the result:

Theorem 5.10

$$F_1 = \frac{1}{12} \ln[\tau_B((a_i)_i)] + \frac{1}{24} \ln \left(\prod_i y'(a_i) \right)$$

τ_B is the Bergman tau function of our problem, which is a function of the position of the branch points in the x -plane, $(a_i)_i$, and which is defined by:

$$\forall i, \quad \frac{\partial \ln \tau_B}{\partial a_i} = S_B(u(a_i))$$

The very special expression of $\omega_3^{(0)}$ (Thm. 4.3, which is a sort of WDVV formula) ensures that the form $\sum_i da_i S_B(u(a_i))$ is closed.

6 Large maps

As an application of the topological recursion, we can derive the critical exponents for the $\mathcal{O}(\mathfrak{n})$ model without loop at boundaries, thus extending the proofs existing for planar maps (genus 0). In this section, we recall known facts about the critical points, outline the method to take the critical limit in the formalism we used, and give the main results. We recall that:

$$\mathfrak{n} = -2 \cos(\pi\mu), \quad \text{with } \mu \in]0, 1[$$

6.1 Principles

6.1.1 Asymptotic of maps

Let $\hat{V}$ be a fixed potential. The analysis of singularities (also called critical points) in t of $W_k^{(g)}(t)$ allows to find the asymptotic of the (weighted) number of $\mathcal{O}(\mathbf{n})$ maps with genus g and k boundaries. We illustrate this on $F_0(t)$.

$$F_0(t) = \sum_{v \geq 0} F_{0|v} t^v$$

was defined as a generating function of spherical $\mathcal{O}(\mathbf{n})$ maps with v vertices. It is a formal power series in t , and we know (see next paragraph) by a corollary of the 1-cut lemma that its radius of convergence $\rho_0(\hat{V})$ is strictly positive. Hence, there exists $t^* \in \mathcal{C}(0; \rho_0(\hat{V}))$ such that $F_0(t)$ has a singularity when $t \rightarrow t^*$. We can decompose $F_0(t) = f_r(t) + f_s(t)$ with f_r analytic in a neighborhood of t^* and f_s nonanalytic (maybe divergent) at $t = t^*$. Then, we have the well-known relation:

$$\begin{aligned} & \text{if } f_s(t) \sim A \left(1 - \frac{t}{t^*}\right)^\alpha && \text{when } t \rightarrow t^* \\ & \text{then } F_{0|v} \sim \frac{A}{\Gamma(-\alpha)} v^{-(\alpha+1)} (t^*)^{-v} && \text{when } v \rightarrow \infty \end{aligned} \quad (6-1)$$

6.1.2 Critical points of the $\mathcal{O}(\mathbf{n})$ model

We have to find the singularities of the $W_k^{(g)}$'s which are the closest to 0 in order to study observables on large maps. We claim that this singularity is common to k, g . Indeed, according to the 1-cut lemma, $W_1^{(0)}(t)$ has a strictly positive radius $\rho_0(\hat{V})$ and a singularity at some $t^* \in \mathcal{C}(0; \rho_0(\hat{V}))$. Successive applications of the loop insertion operator $\frac{\partial}{\partial \hat{V}(x)}$ preserve the radius and the singularity, and yield $W_k^{(0)}$ ($k \geq 1$). We have to argue that the residue formula do not change the radius either, but this granted, t^* is common to all stable $W_k^{(g)}$'s. The unstable functions, F_0 and F_1 are obtained by an integration formula, which preserves the radius and the singularity at t^* .

In general, when one solves the saddle point equation, one finds $y(x) \sim 2\eta_i \sqrt{x - a_i}$ for some $\eta_i \in \mathbb{C}$ when $x \rightarrow a_i$. This means that, in the local parameter on the spectral curve $u \propto \sqrt{x - a_i}$, y has a simple zero at $u = u(a_i)$. This property falls down for some exceptional potentials at some value $t = t^*$, and we rather have $y(x) \propto (x - a_i)^\lambda$ for $\lambda \geq \frac{1}{2}$. Then, the stable $W_k^{(g)}$'s expressed by the residue formula diverge when $t \rightarrow t^*$. The intuition can be supported by the Coulomb gas picture and one finds [32, 14] different kinds of critical points t^* .

- Pure gravity. $a_i^2(t^*)$ is a zero of order p of $A(x^2)$ and $B(x^2)$ in Eqn. 3-1. Then, $y(x) \propto (x - a_i)^{p+\frac{1}{2}}$ when $x \rightarrow a_i$. It happens when the liquid of eigenvalues (in the

thermodynamic limit) crosses a potential barrier, i.e. at a point a_i where $y(a_i) = 0$ and where the effective potential for an eigenvalues, $V_{\text{eff}}(x) = \frac{2V'(x) - nV'(-x)}{4-n^2}$, satisfies $V_{\text{eff}}^{(l)}(a_i) = 0$ for $1 \leq l \leq p+1$. Then, the model has a well-defined limit for x close to a_i , which does not feel the presence of the interface at $x = 0$. Back to combinatorics, such a singularity is associated with large maps without macroscopic loops. This limit exists already in the limit of the 1-matrix model and describes pure gravity. In the CFT classification, it is described by the $(p, 2)$ minimal model.

- $a(t^*) = 0$. The cuts $[-b, -a]$, $[a, b]$ merge. It happens when the eigenvalues can be as close as they want to their mirrors. Then, the method of Section 2 is not valid *stricto sensu*, because the torus is pinched at the point $x = 0$. The singularity lie in D_μ , and not in the polynomials A and B . These critical points are specific to the $\mathcal{O}(n)$ model and associated to large maps where macroscopic loops are densely drawn.
- Combinations of the two situations. Merging cuts ($a = 0$) and tuning of V such that $A(x^2)$ and $B(x^2)$ have zeroes of order m . These multicritical points are also associated to large maps where loops are macroscopic. Although, they are not dense : we rather have cohabitation of regions dominated by gravity, and regions dominated by macroscopic loops [32].

We let aside pure gravity, and concentrate ourselves on the critical point $t \rightarrow t^*$ corresponding to $a \rightarrow 0$.

6.1.3 Taking the limit $a \rightarrow 0$

When $a \rightarrow 0$:

$$K \sim \frac{\pi}{2}, \quad K' \simeq \ln \left(\frac{4b}{a} \right) \rightarrow \infty, \quad \tau = \frac{iK}{K'}$$

The limit of the functions involved depends on the position of x in the complex plane (of u in the torus). We are first interested in the vicinity of a , i.e. $x/a = x^*$ finite. We may use the parametrization:

$$x^* = \sin \varphi = \text{ch } \chi \tag{6-2}$$

When x is in the first quadrant of the physical sheet, $\sigma(x) = a^2 \cos \varphi$, and the cut is located at $\chi \in [0, \infty]$. To take the limit $\tau \rightarrow 0$ in theta functions, it is convenient to perform the modular transformation $\tau \mapsto \tau' = -1/\tau$. Then, in the sum:

$$\vartheta_1(w|\tau') = i \sum_{m \in \mathbb{Z}} (-1)^m e^{i\pi(m-\frac{1}{2})^2 \tau' + i\pi(2m-1)w},$$

only one or two terms (depending on the position of w we are interested in) are dominant since $i\pi\tau' \rightarrow -\infty$. For instance, the special solution of Eqn. 2-4 we introduced in Eqn. 2-5 becomes:

$$D_\mu(u) = \frac{x(u)}{\sigma(x(u))} \frac{\vartheta_1\left(\frac{\varphi}{2K} - \frac{\mu\tau'}{2} \middle| \tau'\right)}{\vartheta_1\left(\frac{\varphi}{2K} \middle| \tau'\right)} \frac{\vartheta_1\left(\frac{\tau'}{2} \middle| \tau'\right)}{\vartheta_1\left((1-\mu)\frac{\tau'}{2} \middle| \tau'\right)} \exp\left[i\pi\mu\left(-\frac{\varphi}{2K} + \frac{\tau'}{2}\right)\right]$$

where $u = \frac{i\varphi}{2K'} + \frac{\tau}{2}$. Taking the limit, we obtain

$$D_\mu(x) \sim \left(\frac{a}{4b}\right)^{-\mu} \frac{e^{-i\mu\varphi}}{2ib \cos \varphi} \quad (6-3)$$

The function for x arbitrary is given by analytic continuation of the expression above. With similar methods, we can derive (see also [14, 18]):

- (see [14, 18] or Appendix D):

$$\alpha_1 \sim -i\mu b, \quad e_\mu^2 \sim -\frac{1}{4} a^{2(1-\mu)} (4b)^{2\mu}$$

The behavior of e_μ can be derived by computing $D_\mu(x)D_\mu(-x)$ computed directly with Eqn. 6-3, and Eqn. 4-3 in Appendix D.

- Basis of 1-cut solution of Eqn. 2-4.

$$\frac{f_\mu}{R_\mu} \sim \text{cte} \cos \varphi \cos \mu \left(\varphi - \frac{\pi}{2}\right) = -i \text{cte} \text{ch} \chi \text{sh} \mu \chi \quad (6-4)$$

$$\frac{\widehat{f}_\mu}{\widehat{R}_\mu} \sim \widehat{\text{cte}} \sin \varphi \sin \mu \left(\varphi - \frac{\pi}{2}\right) = i \widehat{\text{cte}} \text{ch} \chi \text{sh} \mu \chi \quad (6-5)$$

- Basic blocks of the spectral curve (Section 3.2)

$$\begin{aligned} y_\mu &\sim \text{cte} \sin \varphi \cos \mu \left(\varphi + \frac{\pi}{2}\right) = \text{cte}' \text{ch} \chi \text{ch} \mu (\chi + i\pi) \\ \hat{y}_\mu &\sim \widehat{\text{cte}}' \cos \varphi \sin \mu \left(\varphi + \frac{\pi}{2}\right) = -i \widehat{\text{cte}}' \text{sh} \chi \text{sh} \mu (\chi + i\pi) \end{aligned}$$

- Brick of the Cauchy kernel (Thm. 3.3).

$$H_+(\chi_0, \chi) \sim e^{-i\pi\mu} \frac{d\chi_0}{dx^*(\chi_0)} e^{\mu(\chi+\chi_0)} (-1 - i \coth(\chi + \chi_0)) \quad (6-6)$$

- Bergman kernel (Section 3.4).

$$\mathcal{B}(\chi_0, \chi) \sim d\chi_0 d\chi e^{\mu(\chi+\chi_0)} \left[\mu + i\mu \coth(\chi + \chi_0) - \frac{1}{\text{sh}^2(\chi + \chi_0)} \right] \quad (6-7)$$

At the critical point : $a = 0$	
λ	$y(x) \sim C x^\lambda \quad \text{when } x \rightarrow 0$

Approach of the critical point : $a \rightarrow 0$	
α	$\mathcal{L}_{a,b} : y(x) \sim a^\alpha y^*(x/a) \quad x/a \text{ finite}$
$\alpha(k, g)$	$W_k^{(g)}[\mathcal{L}_{a,b}](x_1, \dots, x_k) \sim a^{\alpha(k,g)} W_k^{(g)}[\mathcal{L}^*]\left(\frac{x_1}{a}, \dots, \frac{x_k}{a}\right) \quad 2g - 2 + k \geq 0$
β	$\left(1 - \frac{t}{t^*}\right) \propto a^{2\beta}$
γ_{str}	$U(t) = \frac{\partial^2 F_0}{\partial t^2} \propto \left(1 - \frac{t}{t^*}\right)^{-\gamma_{str}}$

6.2 Spectral curve in the scaling limit

Description of the critical point

We gathered above the definitions of the critical exponents of the bulk $\mathcal{O}(\mathbf{n})$ model. $U(t)$ is a fundamental quantity to many aspects. In the correspondence of the critical model with a CFT, the "string susceptibility" γ_{str} gives the necessary central charge:

$$c = 1 - 6 \frac{\gamma_{str}^2}{1 - \gamma_{str}}$$

Furthermore, the conformal dimensions (from which the scaling exponents can be extracted) of an operator in a CFT coupled to gravitation ($\tilde{\Delta}$), or not (Δ), are related by the KPZ equation [38]:

$$\tilde{\Delta} = \frac{\Delta(\Delta - \gamma_{str})}{1 - \gamma_{str}}$$

In other words, this equation relates exponents of a statistical model of a regular lattice to those on the random lattice.

Limit curve

An important feature of the residue formula is its compatibility with limits of curves. So, we ask ourselves what is the spectral curve $\mathcal{L}_{a,b}$ in the limit $a \rightarrow 0$, and we call it $\mathcal{L}^*$. We find, by linear combination of Eqn. 6-4 and 6-5, that $w_0(\chi) = \text{ch}(\mu + 1)\chi$ and $w_{-1}(\chi) = \text{ch}(\mu - 1)\chi$ are solutions. By choosing appropriate polynomials $A(\text{ch}^2 \chi)$

and $B(\text{ch}^2\chi)$ of maximal degree D , which are also linear combinations of $\text{ch}(2m\chi)$ for $0 \leq l \leq D$ integer, we find recursively that the general solution for w is a linear combination of:

$$w_m(\chi) = \text{ch}(\mu + 2m + 1)\chi, \quad \text{for } -(D + 1) \leq m \leq D$$

The spectral curve associated to a homogeneous part of the resolvent which would be w_m is:

$$y_m(\chi) = 2w_m(\chi) + \mathbf{n}w_m(-(\chi + i\pi)) = -2 \text{sh}[(\mu + 2m + 1)\chi + 2i\pi\mu]$$

We obtain:

Theorem 6.1 *When $a \rightarrow 0$ and $\mathbf{x}^* = \mathbf{x}/a$ is kept finite, the rescaled limit of the spectral curve is of the form:*

$$\begin{cases} x^*(\chi) = \text{ch}\chi \\ y^*(\chi) = \sum_{m=-(D+1)}^D c_m \text{sh}((\mu + 2m + 1)\chi + 2i\pi\mu) \end{cases}$$

If the potential V has maximal degree $d_{\max}$, then $D = \lfloor (d_{\max} - 1)/2 \rfloor$ and $c_m \in \mathbb{C}$. The coefficients are subjected to the extra condition $y(\chi = 0) = 0$.

Eventually, we gather in Fig. 8 the remaining data which initialize the topological recursion in the scaling limit $\mathbf{x}/a$ finite, $a \rightarrow 0$.

Double scaling limit	
$\mathcal{L}^*$	$\mathcal{L}^* : \begin{cases} x^*(\chi) = \text{ch}\chi \\ y^*(\chi) = \sum_{m=-(D+1)}^D c_m \text{sh}((\mu + 2m + 1)\chi + 2i\pi\mu) \end{cases}$
H_+	$H_+^*(\chi_0, \chi) = e^{-i\pi\mu} \frac{d\chi_0}{dx^*(\chi_0)} e^{\mu(\chi+\chi_0)} (-1 - i \coth(\chi + \chi_0))$
$\mathcal{B}^*$	$\mathcal{B}^*(\chi_0, \chi) = d\chi_0 d\chi e^{\mu(\chi+\chi_0)} \left[\mu + i\mu \coth(\chi + \chi_0) - \frac{1}{\text{sh}^2(\chi + \chi_0)} \right]$
$\mathcal{K}^*$	$\mathcal{K}^*(\chi_0, \chi) = \frac{1}{\sin \pi\mu} \frac{d\chi_0}{[y^* dx^*](\chi)} \left[\text{ch} \mu(\chi - i\pi) \text{ch} \mu \chi_0 - \frac{i}{2} \frac{\text{sh} \mu(\chi + \chi_0 - i\pi)}{\text{sh}(\chi + \chi_0)} \text{ch}(\chi + \chi_0) - \frac{i}{2} \frac{\text{sh} \mu(\chi - \chi_0 - i\pi)}{\text{sh}(\chi - \chi_0)} \text{ch}(\chi - \chi_0) \right]$

Figure 8: Summary on the critical $\mathcal{O}(\mathbf{n})$ model.

6.3 Topological recursion in the scaling limit

We see on Eqn. 6-7 that $\mathcal{B}$ has a well-defined limit, without scaling factor, when $a \rightarrow 0$:

$$dx_0 dx \overline{W}_2^{(0)}(x_0, x) \sim dx_0^* dx^* (\overline{W}_2^{(0)})^*(x_0^*, x^*)$$

Subsequently, for the recursion kernel:

$$K(x_0, x) \sim a^{-(\alpha+1)} K^*(x_0^*, x^*)$$

For g, k such that $2g - 2 + k > 0$, $W_k^{(g)}$ is obtained by a stack of $2g - 2 + k$ residues against a recursion kernel, of a product of $\overline{W}_2^{(0)}(\cdot, \cdot)$ blocks. Hence:

$$dx_1 \cdots dx_k W_k^{(g)}(x_1, \dots, x_k) \sim a^{(\alpha+1)(2-2g-k)} dx_1^* \cdots dx_k^* (W_k^{(g)})^*(x_1^*, \dots, x_k^*)$$

This yields:

$$\boxed{\alpha(k, g) = (2 - 2g - k)(\alpha + 1) - k} \quad (6-8)$$

Thus, we have proved the KPZ scaling [27, 8, 9, 38].

6.4 Critical exponents

Determination of λ

We have to solve the saddle point equation for $a = 0$ and x finite. A first method reviewed in Appendix E consists in a direct guess of a good parametrization or/and of the general solution [14, 23, 24]. A second method is to compute directly the limits of theta functions, of D_μ and then of the basis $(f_\mu/R_\mu, \widehat{f}_\mu/\widehat{R}_\mu)$, as we did when x was of order a . Of course, the two methods lead to the same results.

Theorem 6.2 *We introduce the parametrization $x = \frac{b}{\text{ch}\zeta}$. Then, the general spectral curve when $a = 0$ and x is kept finite, is of the form:*

$$\boxed{y(x) = A(x^2) \text{th}\zeta \text{ch}(\mu\zeta) + B(x^2) \text{sh}(\mu\zeta)} \quad (6-9)$$

where A and B are polynomials. If the potential V has maximal degree $d_{\max}$, we have:

$$\deg(A) \leq \left\lfloor \frac{d_{\max}}{2} \right\rfloor - 1, \quad \deg(B) \leq \left\lfloor \frac{d_{\max} - 1}{2} \right\rfloor$$

□

	Table of exponents	Example : triangular lattice	
	Phase $(\varepsilon, m) \in \{\pm 1\} \times \mathbb{N}$	Dense phase $(\varepsilon, m) = (-1, 0)$	Dilute phase $(\varepsilon, m) = (1, 0)$
λ	$\varepsilon(1 - \mu) + 2m + 1$	μ	$2 - \mu$
α	$\varepsilon(1 - \mu) + 2m + 1$	μ	$2 - \mu$
$\alpha(k, g)$	$(2 - 2g)(\alpha + 1) - k\alpha$		
β	$\frac{1-\varepsilon}{2}\mu + m + \frac{1+\varepsilon}{2}$	μ	1
γ_{str}	$\frac{\mu-1}{\frac{1-\varepsilon}{2}\mu+m+\frac{1+\varepsilon}{2}}$	$1 - \frac{1}{\mu}$	$-(1 - \mu)$

Figure 9: Summary on the critical $\mathcal{O}(\mathbf{n})$ model.

When $x \rightarrow 0$, $x \sim \frac{b}{2}e^{-\zeta}$. Since $\mu \in]0, 1[$, the two terms (y_μ and $\hat{y}_\mu$) admit $x^{-\mu}$ as leading order, and:

$$x^\mu, x^{-\mu+2}, \dots, x^{(\mu-1)+2m+1}, x^{(1-\mu)+2m+1}, \dots$$

as subleading orders. If we demand⁹ $\lim_{x \rightarrow 0} y(x) = 0$, A and B are such that this leading order disappears. Hence, the first possible term is x^μ , and other subleading orders can be canceled by a special choice of A and B . Thus, in general, the leading term is x^λ , with:

$$\lambda = \varepsilon(1 - \mu) + 2m + 1, \quad \varepsilon \in \{\pm 1\}, m \in \mathbb{N}$$

The admissible values for ε and m depends on $d_{\max}$. They determine the "phases" of the $\mathcal{O}(\mathbf{n})$ model. For example :

- On a triangular lattice ($d_{\max} = 3$), there exists only two phases, in agreement with [24]:

$$\text{Dense phase} \quad (\varepsilon, m) = (-1, 0) \quad \lambda = \mu$$

$$\text{Dilute phase} \quad (\varepsilon, m) = (1, 0) \quad \lambda = 2 - \mu$$

⁹It is true for all $a \neq 0$ that $y_a(a) = 0$. Though, we did not find a convincing argument to rule out from the case $y_{a=0}(0) = \infty$ for combinatorics. In this case, y could be divergent as $x^{-\mu}$ but integrable on $[0, b]$ since $\mu \in]0, 1[$.

- In the fully packed case ($d_{\max} = 2$), only the dense phase $(\varepsilon, m) = (-1, 0)$ is present, and the limit spectral curve is (see Appendix E):

$$y_{\text{FPL}}(x) = \text{cte} (\text{ch}(\mu\zeta) - \mu \coth\zeta \text{sh}(\mu\zeta)) \quad (6-10)$$

- With the general potential of degree $d_{\max} = 2d' + 1$ ($d' > 1$), there are $2d'$ phases described by $(\varepsilon, m) \in \{\pm 1\} \times [1, d']$.
- With the general potential of degree $d_{\max} = 2d'$, $(\varepsilon, m) = (1, d')$ cannot be reached, so there are $2d' - 1$ phases.

Determination of α

α is defined such that $y_a(x) \sim a^\alpha y^*(x/a)$ for finite x/a and $a \rightarrow 0$. On the other hand, $y_{a=0}(x)$ behaves as x^λ when $x \rightarrow 0$. The two solutions match if $y^*(x^*) \propto x^{\alpha'}$ when $x^* \rightarrow \infty$ with $\alpha = \alpha'$, and $\alpha' = \lambda$. Thms. 6.1 and 6.2 are compatible in the sense that they produce the same possible values for α' and λ .

Determination of β

We have proved that $\left(\partial_t W_1^{(0)}\right)(x) = f_\mu(x)$. Independently, we have in the limit $a \rightarrow 0$:

$$\begin{aligned} \partial_t(W_1^{(0)})(x) &\propto a^{\alpha-2\beta} \times (\text{function of } x/a) && \text{almost by definition} \\ f_\mu(x) &\sim a^{-\mu} f_\mu^*(x/a) && \text{from Eqn. 6-3} \end{aligned}$$

Hence:

$$2\beta = \alpha + \mu$$

Determination of γ_{str}

We take the limit $a \rightarrow 0$ in Eqn. 5-7 giving the third derivative of F_0 and keep the nonanalytic part at $t = t^*$:

$$\left(\frac{\partial^3 F_0}{\partial t^3}\right)_{\text{singular}} \sim \text{cte } a^{2(1-\mu)-2\beta} \quad (6-11)$$

Hence:

$$\gamma_{str} = \frac{\mu - 1}{\beta} = \frac{\mu - 1}{\frac{1+\varepsilon}{2}\mu + m}$$

(6-12)

We notice that γ_{str} is always negative. Furthermore, the limit of Eqn. 5-7 yields a regular part for $\partial_t^3 F_0$, i.e which is analytic at $t = t^*$. This regular part is dominant in $\partial_t^3 F_0$ when $t \rightarrow t^*$. Also, the same phenomenon occurs for $U(t)$ and $F_0(t)$.

Critical behavior of F_1

With expression Eqn. 5-9, it is easy to see that $F_1 \propto \ln(a)$ when $a \rightarrow 0$.

6.5 Remark: double scaling limit

We have reviewed the fact that, for a given value of $\mathfrak{n}$, the model has many possible continuum limits, described by $(\varepsilon, m) \in \{\pm 1\} \times \mathbb{N}$. We refer to [14] or [10] for a discussion of the case " $\mu = p/q$ rational" in relation with the (p, q) minimal models of CFT.

For any of these limits, we have seen that the stable $W_k^{(g)}(t)$ diverge (at least for $g \geq 2$) as $a^{\alpha(k,1)-(2g-2)(\alpha+1)}$ when $t \rightarrow t^*$. Take $k \neq 0$ to avoid unstable maps, and consider the formal power series $W_k(t)$, depending on N :

$$\begin{aligned} W_k(t|N) &= \sum_{g \geq 0} \left(\frac{N}{t} \right)^{2-2g} W_k^{(g)}(t) \\ &= a^{-\alpha(k,1)} \sum_{g \geq 0} \left(\frac{Na^{-(\alpha+1)}}{t} \right)^{2-2g} a^{\alpha(k,g)} W_k^{(g)}(t) \end{aligned}$$

We can also define a function:

$$W_k^*(N^*) = \sum_{g \neq 0} (N^*)^{2-2g} W_k^{(g)*}$$

If we send $a \rightarrow 0$ and $N \rightarrow \infty$ while keeping $Na^{-(\alpha+1)} = N^*$ finite, then:

$$W_k(t|N) \sim W_k^*(N^*)$$

For this reason, W_k^* , and by extension $W_k^{(g)*}$, are called in matrix model context the "double scaling limit" of W_k , resp. $W_k^{(g)}$. Within the conjecture relating limits of matrix models to CFT (see Section 1.4), W_k^* should be solution of PDE's coming from the conformal field theory of central charge:

$$c = 1 - 6 \left(\sqrt{\lambda} - \frac{1}{\sqrt{\lambda}} \right)$$

Still, it demands a mathematical proof.

7 Conclusion

We have defined for the first time the topological recursion for a spectral curve which has monodromies around branchpoints, and extended to that case all its usual properties. The bulk correlation functions $W_k^{(g)}$ in the $\mathcal{O}(\mathfrak{n})$ matrix model fit in the formalism.

So, we have an algorithm to compute number of maps with self avoiding loops of $\mathbf{n}$ possible colors in any topology and with an arbitrary number of boundaries. In particular, the generating function of cylinders $W_2^{(0)}$ is universal (as expected) and closely related to a deformation $\wp_\mu$ of the Weierstraß $\wp$ function. We also derived new formulas for F_0 and F_1 . Our method consisted in finding a good description of the ring of meromorphic functions on the spectral curve. The theory of elliptic functions can be generalized to functions periodic in 1 direction, and taking a phase in the other direction. $\wp_\mu$ give rise to a Bergman kernel (fundamental bi-differential with a double pole at coinciding points), which can be used to express all meromorphic forms on the spectral curve.

The generalization to the multi-cut solutions of the $\mathcal{O}(\mathbf{n})$ model loop equations seems straightforward: in the residue formula expressing $W_k^{(g)}$, one just has to pick up a residue at every endpoint of a cut. Some work remains however. We must first express the general multicut solution to the master loop equation (Thm. ??), and find the appropriate Bergman kernel. We will return to this point in the future. We stress that for combinatorics of maps, the 1-cut solution that we presented is enough.

The next step is to compute all correlations functions with residue formulas, in the spirit of what was done for the 2-matrix model [21]. It would be very interesting to explore the integrability of the model before the continuum limit, and then, the relations with boundary conformal field theory. Since we can compute a lot of observables in the discrete model, we think that many exact results for the continuum limit could be obtained by this method. For instance, we have been able to give a rigorous proof of the KPZ scaling for all topologies.

On the flat lattice, there exists a relation between the $\mathcal{O}(\mathbf{n})$ model, and the Potts model with $\mathbf{q} = \mathbf{n}^2$ colors. Namely, the critical point of the $\mathbf{n}^2$ Potts model on a square lattice is equivalent to the critical point of the $\mathcal{O}(\mathbf{n})$ model on a square-triangular lattice in the dense phase. On the random lattice, there is again a relation between the two models. They both admit a matrix model representation, and the planar resolvents $W_1^{(0)}(x)$ of these two matrix models are (up to some simple change of variable) reciprocal functions [3]. It would be interesting to see if a topological recursion could solve the $\mathbf{q}$ -Potts matrix model for all topologies, and then, if a relation with the $\mathcal{O}(\mathbf{n})$ model can be extended to all topologies. The Potts model has been up to now more popular in combinatorics and statistical physics than the $\mathcal{O}(\mathbf{n})$ model. Nevertheless, supported by the present work, we think that the $\mathcal{O}(\mathbf{n})$ on the random lattice is easier to solve, and that this solution might shed light on the Potts model on random lattice (outside and at the critical point).

We also expect that, for many other models for which the leading order resolvent $W_1^{(0)}$ is known, like the $\widehat{A}, \widehat{D}, \widehat{E}$ models of [33], or the $ABAB$ model considered in

[42], a topological recursion (maybe slightly deformed) can be found to find all genus correlation functions. We are currently working on this problem.

Acknowledgments

We thank I. Kostov and J.E. Bourgine for valuable discussions and indication of references, and also J. Bouttier and N. Curien.

A Proof of the 1-cut lemma

Lemma A.1 *For all $\mathbf{n}$, k , g , there exists $\rho > 0$ such that $\#\mathbb{M}_k^{(g)}(v) \in O(\rho^v)$ when $v \rightarrow \infty$.*

proof:

This is based on a rough upper bound on the cardinality of $\mathbb{M}_k^{(g)}$. If $\mathbf{n} = 0$ and $t_j = 0$ for $j \geq 4$, $\mathbb{M}_k^{(g)}(v)$ are sets of usual triangulations, and it is known [40] that the lemma is true with $\bar{\rho} = \frac{1}{6\sqrt{3}}$. We assume $\mathbf{n}$ an integer, for it is enough to prove the lemma for $[\mathbf{n}]$. Now, we describe in Fig. A a map from $\mathbb{M}_k^{(g)}(v)$ to the set of triangulations with v vertices where every face carries a label among $\{\text{B}, \text{I}, \text{M}, 1, \dots, \mathbf{n}\}$.

This association is injective:

- If a decorated triangulation comes from $\mathcal{M} \in \mathbb{M}_k^{(g)}(v)$, the unmarked edges in $\mathcal{M}$ are those which glue two "B"-triangles, while the marked edges glue a "B" and a "M" triangle. So, we have the skeleton of $\mathcal{M}$.
- The polygons of this skeleton admitting an inner triangle labeled $j \in \{1, \dots, \mathbf{n}\}$ instead of "I" are necessarily triangles. They must carry a loop, and the j -label indicates where it should be drawn.

By construction:

$$v(\mathcal{M}') \leq (d_{\max} + 1)v(\mathcal{M}), \quad g(\mathcal{M}) = g(\mathcal{M}'), \quad k(\mathcal{M}) = k(\mathcal{M}')$$

Hence:

$$\#\mathbb{M}_k^{(g)}(v) \in O(\bar{\rho}^{(\mathbf{n}+3)(d_{\max}+1)v})$$

□

We proceed in detail with the proof of the 1-cut lemma, which is now the same as for usual maps and the 1-matrix model. We recall that the notion of the color of a loop does not intervene in $\mathbb{M}_k^{(g)}$, but only in the weight of a map: the previous upper bound on $\#\mathbb{M}_k^{(g)}$ is valid uniformly for $\mathbf{n}$ bounded. Moreover, we notice that the association of Fig. A preserves the number of automorphism. We conclude that the coefficient of t^v is a $O(\rho_0/r)$ where

$$\begin{aligned} \rho_0 &= \max(t_j^{\frac{2}{j-2}}, \mathbf{n}c^2) \\ r &= \min_i \sqrt{\left|x_i - \frac{c}{2}\right|} \end{aligned}$$

which proves the first part of the 1-cut lemma.




Figure 10: Every j -gon (with or without loops, marked or unmarked) is itself triangulated with $3j$ decorated triangles: construct a triangle on each edge, and a triangle at each corner. In the inside, there remains a j -gon which you cut in j triangular sectors from its center, that you label "I". The other labels are distributed as follows: for a triangle carrying a loop, the corner triangle where the loop is drawn is decorated by the color of the loop $j \in \{1, \dots, \mathbf{n}\}$; for a marked face, the triangle constructed on the marked edge is decorated by "M" ; in any other case, the triangles constructed on the edges are decorated by "B", while the other are labelled "I".

Next, we solve the master loop equation in $W_1^{(0)}(x)$ which is quadratic:

$$\begin{aligned} W_1^{(0)}(x) &= \frac{1}{2} \left(V'(x) - \mathbf{n}W_1^{(0)}(-x) + \sqrt{\Delta(x)} \right) \\ \Delta(x) &= \left(V'(x) - \mathbf{n}W_1^{(0)}(-x) \right)^2 \\ &\quad - 4 \left(P_1^{(0)}(x) + P_1^{(0)}(-x) - V'(-x)W_1^{(0)}(-x) + W_1^{(0)}(-x)^2 \right) \end{aligned}$$

We have to describe the zeroes of Δ for t small enough. There exists r small enough such that the discs of centers $\pm \frac{c}{2}$ and radius r do not intersect and such that V' has no other zero than $\frac{c}{2}$ in $\mathcal{D}(\frac{c}{2}; r)$. Then, for $t \in \mathcal{D}(0; \frac{\rho_0}{r})$, we see that $\Delta(x)$ is holomorphic on $\mathcal{D}(\frac{c}{2}; r)$. For $t = 0$, $\Delta(x) = (V'(x))^2$ is not the zero function and has a double zero in c . By continuity in t and analyticity for $x \in \mathcal{D}(\frac{c}{2}; r)$, Δ has still two zeroes (with multiplicity), which limit is $\frac{c}{2}$ when $t \rightarrow 0$. Thus, there exists an holomorphic function h_t in $\mathcal{D}(\frac{c}{2}; r)$, without zeroes, and a continuous pair $\{a(t), b(t)\}$ such that $\Delta(x) = h_t^2(x)(x - a(t))(x - b(t))$.

Computing $\Delta(a_i(t))$ for a zero $a_i(t)$ of Δ from Eqn. 1-1 allows us to find $a_i(t)$ as a formal series in $\sqrt{t}$, and yields two solutions $a(t)$ and $b(t)$, such that $a(t) + b(t) \in O(t)$. The property of having a finite radius of convergence in t for $W_1^{(0)}$ implies a finite radius of convergence for a and b .

Eventually, the analytical properties of $W_k^{(g)}$ are derived by a straightforward recursion using the loop equations and Lemma 2.1.

B Elliptic functions

Basic facts

Let us say a few words about bi-periodic functions, i.e. satisfying:

$$f(w) = f(w + 1) = f(w + \tau) \tag{2-1}$$

where $(1, \tau)$ is $\mathbb{R}$ -free. Such functions are also called "elliptic functions" [25]. They satisfy a few properties:

- If f has no pole, then f is constant.
- The number of zeroes of f in the rectangle $(0, 1, 1 + \tau, \tau)$ equals the number of its poles.
- The sum of residues of f at its poles in the rectangle, vanishes. In particular, this shows that f cannot have only one simple pole, it must have at least two simple poles, or one multiple pole.

ϑ_1 function

Let $\tau \in \mathbb{C}$ such that $\text{Im}\tau > 0$. The first Jacobi theta function is defined by:

$$\vartheta_1(w|\tau) = i \sum_{m \in \mathbb{Z}} (-1)^m e^{i\pi(m-\frac{1}{2})^2 \tau + i\pi(2m-1)w} \quad (2-2)$$

This series is absolutely convergent for all $w \in \mathbb{C}$, so $\vartheta_1(\cdot|\tau)$ is an entire function. It satisfies:

$$\vartheta_1(w|\tau) = -\vartheta_1(w+1|\tau), \quad \vartheta_1(w+\tau|\tau) = e^{-2i\pi(w+\frac{\tau}{2})} \quad (2-3)$$

and:

$$\vartheta_1(w|\tau) = -\vartheta_1(-w|\tau) \quad (2-4)$$

It has a unique zero mod $(\mathbb{Z} \oplus \tau\mathbb{Z})$, located at 0. Besides:

$$(\vartheta_1)'(1/2|\tau) = 0 \quad (2-5)$$

Description of the meromorphic functions on the torus

An elliptical function f with poles $w_1, \dots, w_L$ and poles $w'_1, \dots, w'_{L'}$, can always be written¹⁰ as:

$$f(w) = A \frac{\prod_{l'=1}^{L'} \vartheta_1(w - w_{l'}|\tau)}{\prod_{l=1}^L \vartheta_1(w - w_l|\tau)} \quad (2-6)$$

for some constant A , and the poles and zeroes are such that:

$$\sum_{l'} w'_{l'} - \sum_l w_l = 0 \mod \mathbb{Z} \oplus \tau\mathbb{Z} \quad (2-7)$$

Inversion of modulus

There is a relation between $\vartheta_1(\cdot|\tau)$ and $\vartheta_1(\cdot|\tau')$, where $\tau' = -1/\tau$, which is an application of Poisson's summation formula:

$$\vartheta_1(w|\tau) = \frac{-1}{\sqrt{i\tau}} e^{\frac{-i\pi w^2}{\tau}} \vartheta_1\left(\frac{w}{\tau} \middle| -\frac{1}{\tau}\right) \quad (2-8)$$

Weierstraß function

Let us introduce the Weierstraß $\wp$ function:

$$\begin{aligned} \wp(w|\tau) &= \frac{1}{w^2} + \sum_{(l,m) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \left(\frac{1}{(w+l+m\tau)^2} - \frac{1}{(l+m\tau)^2} \right) \\ &= \sum_{m \in \mathbb{Z}} \frac{\pi^2}{\sin^2 \pi(w+m\tau)} - \sum_{m \in \mathbb{Z} \setminus \{0\}} \frac{\pi^2}{\sin^2 \pi m\tau} - \frac{\pi^2}{3} \end{aligned} \quad (2-9)$$

¹⁰Indeed, the ratio of f and this expression would be an elliptical function with no pole, i.e a constant.

This function is elliptic, and related to the second derivative of $\ln \vartheta_1$:

$$\wp(w|\tau) = -(\ln \vartheta_1)''(w|\tau) + \tilde{c}_0 \quad (2-10)$$

for some constant $\tilde{c}_0$ depending on τ . It satisfies the differential equations:

$$\wp'^2 = 4\wp^3 - g_2\wp - g_3 \quad (2-11)$$

$$\wp'' = 6\wp^2 - \frac{g_2}{2} \quad (2-12)$$

where g_2 and g_3 are two constants depending on τ .

C Properties of the μ -deformed Weierstraß function

Definition and basic facts

We have defined:

$$\wp_\mu(w) = \sum_{m \in \mathbb{Z}} e^{-i\pi(1-\mu)m} \frac{\pi^2}{\sin^2 \pi(w + m\tau)} \quad (3-1)$$

We shall derive the differential equations it satisfies for $\mu \neq 1 \pmod{2\mathbb{Z}}$. Let us recall two essential facts about functions which, like $\wp_\mu$, belong to $\text{Ker}(\mathbf{T}_1 - \text{id}) \cap \text{Ker}(\mathbf{T} - e^{i\pi\nu}\text{id})$.

(a) Any quotient of them is elliptic, and any elliptic function is a rational function of $\wp$ and $\wp'$, modulo $\wp'^2 = (4\wp^3 - g_2\wp - g_3)$.

(b) Any such function has at least a pole and a zero (just represent it by the accurate ratio and products of theta functions).

(a) could be used directly to find $\wp'_\mu/\wp_\mu$ and $\wp''_\mu/\wp_\mu$ by studying their zeroes and poles. We found easier to take another route.

First order equation

For $\mu = 1$, $\wp_\mu$ reduces to the Weierstrass function up to a constant depending on τ :

$$\wp_1(w) = \wp(w) + c_0, \quad c_0 = \frac{\pi^2}{3} + \sum_{m \in \mathbb{Z} \setminus \{0\}} \frac{\pi^2}{\sin^2 \pi m\tau} \quad (3-2)$$

From Eqns. 2-11-2-12:

$$(\wp'_1)^2 = 4\wp_1^3 - 12c_0\wp_1^2 + (12c_0^2 - g_2)\wp_1 - (g_3 + 4c_0^3) \quad (3-3)$$

$$\wp''_1 = 6\wp_1^2 - 12c_0\wp_1 + 6c_0^2 - \frac{g_2}{2} \quad (3-4)$$

where g_2, g_3 are constants depending on τ . In the limit $\mu \rightarrow 1$, we ought to recover these equations. Yet, the differential equation involving only $\wp_\mu$ that we look for must be linear. We guess that the correct generalization of $(\wp'_1(w))^2$ when $\mu \neq 0$ is $\wp'_1(w)\wp'_\mu(w)$, and we try to match its polar part when $w \rightarrow 0$ with a generalization of the RHS of Eqn. 3-3. $\wp_1$ is even and behaves as:

$$\wp_1(w) = \frac{1}{w^2} + c_0 + \frac{1}{2}c_2w^2 + O(w^4) \quad (3-5)$$

where it turns out (from Eqn. 3-3) that $c_2 = g_2/10$. On the other hand, $\wp_\mu$ is not even. So, there exists constants $c_j^{(\mu)}$ ($0 \leq j \leq 3$) such that:

$$\wp_\mu(w) = \frac{1}{w^2} + c_0^{(\mu)} + c_1^{(\mu)}w + \frac{1}{2}c_2^{(\mu)}w^2 + \frac{1}{6}c_3^{(\mu)}w^3 + O(w^4) \quad (3-6)$$

We obtain:

$$\wp'_1(w)\wp'_\mu(w) = \frac{4}{w^6} - \frac{2c_1^{(\mu)}}{w^3} - \frac{2c_2^{(\mu)} + 2c_2}{w^2} + \frac{c_3^{(\mu)}}{w} + O(1) \quad (3-7)$$

The simplest candidate in $\text{Ker}(\mathbf{T}_1 - \text{id}) \cap \text{Ker}(\mathbf{T} - e^{i\pi\nu}\text{id})$ to match this polar part is a linear combination of

$$\wp_1^2\wp_\mu, \quad \wp_1\wp_\mu, \quad \wp_\mu, \quad \wp'_1\wp_\mu, \quad \wp_1\wp'_\mu \quad \text{and} \quad \wp'_\mu. \quad (3-8)$$

The polar part of these six functions are independent, so they are enough to fix the terms up to w^{-6} . As an application of (b), this linear combination must be equal to $\wp'_1\wp'_\mu$. We give directly the result:

$$\wp'_1\wp'_\mu = 4\wp_1^2\wp_\mu - 4(2c_0 + c_0^{(\mu)})\wp_1\wp_\mu + \left[4c_0^2 + c_0c_0^{(\mu)} + (c_0^{(\mu)})^2 - g_2^{(\mu)}\right]\wp_\mu + \lambda q_\mu \quad (3-9)$$

where q_μ is the function defined by:

$$q_\mu(w) = \left[\wp_1(w)\wp'_\mu(w) - \wp'_1(w)\wp_\mu(w) + (c_0^{(\mu)} - c_0)\wp'_\mu(w) - 3c_1^{(\mu)}\wp_\mu(w)\right] \quad (3-10)$$

and:

$$g_2^{(\mu)} \stackrel{\text{def}}{=} 6c_2 + 4c_2^{(\mu)} \quad (3-11)$$

$$\lambda \stackrel{\text{def}}{=} \frac{-\frac{5}{6}c_3^{(\mu)} + (2c_0 + c_0^{(\mu)})c_1^{(\mu)}}{c_2^{(\mu)} - c_2} \quad (3-12)$$

Second order equation

Similarly, we can match the polar part when $w \rightarrow 0$ of $\wp''_\mu$ by a linear combination of

$$\wp_1\wp_\mu, \quad \wp_\mu, \quad \wp'_1\wp_\mu, \quad \wp_1\wp'_\mu \quad \text{and} \quad \wp'_\mu. \quad (3-13)$$

The result is:

$$\wp''_\mu = 6\wp_1\wp_\mu - 6(c_0 + c_0^{(\mu)})\wp_\mu + \lambda'q_\mu(w) \quad (3-14)$$

where:

$$\lambda' = \frac{-3c_1^{(\mu)}}{c_2^{(\mu)} - c_2} \quad (3-15)$$

Consistency of the limit $\mu \rightarrow 1$

Let us consider the limit $\mu \rightarrow 1$ of these differential equations. $c_j^{(\mu)}$ (for $0 \leq j \leq 3$) is equal to c_j (which is zero when j is odd) in the limit $\mu \rightarrow 1$ since we can commute the limit and the residue in:

$$c_j^{(\mu)} = \text{Res}_{w \rightarrow 0} \frac{dw}{w^{l+1}} \wp_\mu(w) \quad (3-16)$$

Comparing to Eqn. 3-9 to 3-3, and Eqn. 3-14 to 3-4, we find:

$$\begin{aligned} \forall w \in \mathbb{C} \quad \lim_{\mu \rightarrow 1} \lambda q_\mu(w) &= -(4c_0^3 + g_3) \\ \lim_{\mu \rightarrow 1} \lambda' q_\mu(w) &= 6c_0^2 - \frac{g_2}{2} \end{aligned} \quad (3-17)$$

But one can compute independently:

$$\lambda' q_\mu(w) = -\frac{6c_1^{(\mu)}}{w} + \lambda' \left(\frac{5}{6}c_3^{(\mu)} - 2c_1^{(\mu)}c_0^{(\mu)} \right) + O(w) \quad (3-18)$$

Hence λ and λ' diverge when $\mu \rightarrow 1$ in a way such that:

$$\lim_{\mu \rightarrow 1} \frac{\lambda}{\lambda'} = \frac{g_3 + 4c_0^2}{\frac{g_2}{2} - 6c_0^3} \quad (3-19)$$

Primitive of $\wp_\mu$

$\wp_\mu$ can be integrated in terms of theta functions. We only state the formula:

$$\int_{w_1}^{w_2} dw \wp_\mu(w) = \pi \frac{\vartheta'_1(0|\tau)}{\vartheta_1(\nu/2|\tau)} \left(\frac{\vartheta_1(\nu/2 - w_1|\tau)}{\vartheta_1(w_1|\tau)} - \frac{\vartheta_1(\nu/2 - w_2|\tau)}{\vartheta_1(w_2|\tau)} \right) \quad (3-20)$$

From this point, one can also obtain theta function representations for $c_j^{(\mu)}$. For example:

$$c_0^{(\mu)} = \frac{1}{6} \frac{\vartheta_1'''(0|\tau)}{\vartheta_1'(0|\tau)} - \frac{1}{2} \frac{\vartheta_1''(\nu/2|\tau)}{\vartheta_1(\nu/2|\tau)} \quad (3-21)$$

$$c_1^{(\mu)} = \frac{1}{6} \left(\frac{\vartheta_1'''(0|\tau)}{\vartheta_1'(0|\tau)} \frac{\vartheta_1'(\nu/2|\tau)}{\vartheta_1(\nu/2|\tau)} - \frac{\vartheta_1'''(\nu/2|\tau)}{\vartheta_1(\nu/2|\tau)} \right) \quad (3-22)$$

D Properties of the special solutions of Eqn. 2-4

- Theta expression for D_μ .

$$D_\mu(u) = \frac{x(u)}{\sigma(x(u))} \frac{\vartheta_1\left(u - \frac{\tau}{2} + \frac{\mu}{2} \middle| \tau\right)}{\vartheta_1\left(u - \frac{\tau}{2} \middle| \tau\right)} \frac{\vartheta_1\left(-\frac{1}{2} \middle| \tau\right)}{\vartheta_1\left(-\frac{1}{2} + \frac{\mu}{2} \middle| \tau\right)} \quad (4-1)$$

$u_e = \frac{\tau - \mu}{2}$ is the second zero (mod $\mathbb{Z} \oplus \tau\mathbb{Z}$) of D_μ . We call $e_\mu = x(u_e)$, the corresponding point in the physical x-sheet:

$$e_\mu = a \operatorname{sn}_k(i\mu K') \quad (4-2)$$

We use the notation $D_\mu(x) = D_\mu(u(x))$.

- Relation between D_μ and $D_{-\mu}$. $D^{(2)} : u \mapsto D_\mu(u)D_\mu(\tau - u)$ is an u -even and bi-periodic function. Hence $D_\mu^{(2)}(x) = D_\mu^{(2)}(u(x))$ is a rational fraction of x^2 . In this variable, it has simple poles at $x^2 = a^2$ and $x^2 = b^2$, simple zeroes at $x^2 = \infty$, $x^2 = e_\mu^2$, and is equivalent to $-1/x^2$ when $x^2 \rightarrow \infty$. Thus:

$$D_\mu^{(2)}(x) = -\frac{x^2 - e_\mu^2}{(x^2 - a^2)(x^2 - b^2)} \quad (4-3)$$

- Asymptotic of $D_\mu(x)$. We define the constants α_1 and α_2 by:

$$D_\mu(x) = \frac{1}{x} + \frac{\alpha_1}{x^2} + \frac{\alpha_2}{x^3} + O\left(\frac{1}{x^4}\right) \quad \text{when } x \rightarrow \infty \text{ in the physical sheet} \quad (4-4)$$

Studying Eqn. 4-1 when $u \rightarrow \frac{-1+\tau}{2}$ yields the expression:

$$\alpha_1 = \frac{-ib}{2K'} (\ln \vartheta_1)' \left(\frac{1-\mu}{2} \middle| \tau \right). \quad (4-5)$$

Studying Eqn. 4-3 when $x \rightarrow \infty$ yields the expression of α_2 :

$$\alpha_2 = \frac{1}{2}(\alpha_1^2 + a^2 + b^2 - e_\mu^2) \quad (4-6)$$

The terms up to α_2 are involved in the fully packed case.

- Derivative of D_μ . D_μ and $(\partial_u D_\mu)$ are both in $\operatorname{Ker}(\mathbf{T} - e^{i\pi\mu})$, so $\partial_u \ln D_\mu$ is 1- and τ - translation invariant. By studying its analytical properties, we obtain:

$$\frac{dD_\mu}{dx} = L(x)D_\mu(x) \quad (4-7)$$

where:

$$L(x) = \frac{-\alpha_1 + I(x)}{\sigma(x)} - \frac{d \ln \sigma}{dx} \quad (4-8)$$

$$\text{and } I(x) = \frac{x\sigma(x) + e_\mu\sigma(e_\mu)}{x^2 - e_\mu^2} \quad (4-9)$$

- Definition of f_μ and $\widehat{f}_\mu$.

$$f_\mu(u) = \frac{D_\mu(u) + D_\mu(-u)}{1 - e^{i\pi\mu}} \quad (4-10)$$

$$\widehat{f}_\mu(u) = \frac{\sigma(x(u))}{x(u)} \frac{D_\mu(u) - D_\mu(-u)}{1 + e^{i\pi\mu}} \quad (4-11)$$

We recall that $u(-x) = \tau - u(x)$. So, in the x variable:

$$f_\mu(x) = \frac{D_\mu(x) + e^{i\pi\mu}D_\mu(-x)}{1 - e^{i\pi\mu}} \quad (4-12)$$

$$\widehat{f}_\mu(x) = \frac{1}{n-2} \frac{\sigma(x)}{x} (nf_\mu(x) + 2f_\mu(-x)) \quad (4-13)$$

- Wronskien. The wronskien function is defined by $\Xi_\mu = (\mathbf{T}f_\mu) \widehat{f}_\mu - f_\mu (\mathbf{T}\widehat{f}_\mu)$. After computation:

$$\begin{aligned} \Xi_\mu(u(x)) &= f_\mu(x)\widehat{f}_\mu(-x) - f_\mu(-x)\widehat{f}_\mu(x) \\ &= -\frac{2\sigma(x)}{x} D_\mu^{(2)}(x) \\ &= \frac{2(x^2 - e_\mu^2)}{x\sigma(x)} \end{aligned} \quad (4-14)$$

- $\perp$ -Norms of f_μ and $\widehat{f}_\mu$. From the definitions:

$$\begin{aligned} R_\mu(x^2) = (f_\mu \perp f_\mu)(x) &= -(2-n) D_\mu^{(2)}(x) \\ &= (2-n) \frac{x^2 - e_\mu^2}{(x^2 - a^2)(x^2 - b^2)} \end{aligned} \quad (4-15)$$

$$\begin{aligned} \widehat{R}_\mu(x^2) = (\widehat{f}_\mu \perp \widehat{f}_\mu)(x) &= -(2+n) \frac{\sigma(x)^2}{x^2} D_\mu^{(2)}(x) \\ &= (2+n) \frac{x^2 - e_\mu^2}{x^2} \end{aligned} \quad (4-16)$$

- Asymptotic of f_μ and $\widehat{f}_\mu$.

$$f_\mu(x) = \frac{1}{x} + \frac{1 + e^{i\pi\mu}}{1 - e^{i\pi\mu}} \frac{\alpha_1}{x^2} + \frac{\alpha_2}{x^3} + O\left(\frac{1}{x^4}\right) \quad (4-17)$$

$$\widehat{f}_\mu(x) = 1 + \frac{1 - e^{i\pi\mu}}{1 + e^{i\pi\mu}} \frac{\alpha_1}{x} + \frac{\alpha_2}{x^2} + O\left(\frac{1}{x^3}\right) \quad (4-18)$$

when $x \rightarrow \infty$ in the physical sheet.

- Values at e_μ . Since $D_\mu(e_\mu) = 0$, $f_\mu(e_\mu)$ and $\widehat{f}_\mu(e_\mu)$ are both proportional to $D_\mu(-e_\mu)$ (everything is read in the variable x). More precisely:

$$\frac{f_\mu(e_\mu)}{\widehat{f}_\mu(e_\mu)} = -\frac{1 + e^{i\pi\mu}}{1 - e^{i\pi\mu}} \frac{e_\mu}{\sigma(e_\mu)} \quad (4-19)$$

- Behavior at branch points. Thanks to Eqn. 4-15, we can find the behavior of f_μ near $a_i \in \{a, b\}$:

$$\lim_{x \rightarrow a_i} \sigma(x) f_\mu(x) = \sqrt{(2 - \mathbf{n})(a_i^2 - e_\mu^2)} \quad (4-20)$$

- First order 2×2 differential system. The differential equation Eqn. 4-7 can be turned into a first order differential system defining $(f_\mu, \widehat{f}_\mu)$ intrinsically. The solution is unique if we require the asymptotic to be given by Eqn. 4-18 up to $O(x^{-3})$ when $x \rightarrow \infty$ in the physical sheet. The data of e_μ (Eqn. 4-2) and α_1 (Eqn. 6-4) ensures that this functions have the correct monodromy, i.e are one cut solutions of the saddle point equation 2-2.

$$\frac{d}{dx} \begin{pmatrix} f_\mu(x) \\ \frac{x \widehat{f}_\mu(x)}{\sigma(x)} \end{pmatrix} = \begin{pmatrix} L_o(x) & \frac{1+e^{i\pi\mu}}{1-e^{i\pi\mu}} L_e(x) \\ \frac{1-e^{i\pi\mu}}{1+e^{i\pi\mu}} L_e(x) & L_o(x) \end{pmatrix} \cdot \begin{pmatrix} f_\mu(x) \\ \frac{x \widehat{f}_\mu(x)}{\sigma(x)} \end{pmatrix} \quad (4-21)$$

where L_o and L_e and the odd and even part of L :

$$L_o(x) = \frac{1}{2} \frac{d \ln R_\mu}{dx} = \frac{d}{dx} \left[\frac{1}{2} \ln(x^2 - e_\mu^2) - \ln \sigma(x) \right] \quad (4-22)$$

$$L_e(x) = \frac{1}{\sigma(x)} \left[-\alpha_1 + \frac{e_\mu \sigma(e_\mu)}{x^2 - e_\mu^2} \right] = \frac{-\alpha_1 x^2 - \widehat{e}_\mu^2}{\sigma(x) x^2 - e_\mu^2} \quad (4-23)$$

We have called $\widehat{e}_\mu = \sqrt{e_\mu^2 + \frac{e_\mu \sigma(e_\mu)}{\alpha_1}}$, the zero of $L_e(x)$.

E Resolvent at $a = 0$ and $b = \infty$

The master loop equation when $a = 0$ (x finite), or $b = \infty$ (equivalently $a \rightarrow 0$ and $x^* = x/a$ finite), can be solved directly, and quite explicitly in terms of Chebyshev functions. These computations were first done by Kostov et al. at the time of the introduction of the $\mathcal{O}(n)$ model. We presented another method to compute when $a \rightarrow 0$ in Section. ??, and original computations with many examples (in particular, the cases μ rational) can be found in [23, 14, 18, 19]. Here, we review in a self-contained way the solution for $b = \infty$, for sake of completeness.

We want to solve the following problem. Let $\gamma = [0, b]$. Find $W = W_1^{(0)}$ such that:

- (i) W is holomorphic on $\mathbb{C} \setminus \gamma$.
- (ii) $W(x) \sim t/x$ when $x \rightarrow \infty$
- (iii) $W(x) \propto \sqrt{x-b}$ when $x \rightarrow b$ (resp
- (iii) For any $x \in \gamma$, we have the linear equation:

$$W(x + i0^+) + W(x + i0^-) + nW(-x) = V'(x) \quad (5-1)$$

Then, the density $\rho(x)$, defined for $x \in \gamma$ is recovered through:

$$\rho(x) = \frac{1}{2i\pi t} (W(x - i0^+) - W(x + i0^+)) \quad (5-2)$$

We also set $n = -2 \cos \pi \mu$, with $\mu \in [0, 1]$. The relation between b (if $\gamma = [0, b]$) or a (if $\gamma = [a, \infty]$), and t , must be such that $\int_0^b \rho = 1$.

At the end, we specialize the results to the fully packed loop model, where the potential is quadratic $V(x) = \frac{1}{2} (x - \frac{c}{2})^2$.

Properties of the linear equation

- The linear equation has an easy particular solution, so we can write:

$$W(x) = \overline{W}(x) + \frac{2V'(x) - nV'(-x)}{4 - n^2} \quad (5-3)$$

with $\overline{W}(x)$ satisfying:

$$\overline{W}(x + i0^+) + \overline{W}(x - i0^+) + n\overline{W}(-x) = 0 \quad (5-4)$$

- Let f and g are meromorphic functions with one cut on γ , which are solution of Eqn. 5-4. Then, the quantity:

$$(f|g)(x) = f(x)g(x) + f(-x)g(-x) + \frac{n}{2} (f(x)g(-x) + f(-x)g(x)) \quad (5-5)$$

is meromorphic, even, and has no cut. Hence, it is a rational function of x^2 .

- The vector space of solutions of Eqn. 5-4, over the field of rational functions of x^2 , has dimension 2.

Chebyshev functions

A basis of functions having similar properties is known, namely the Chebyshev functions $T_\mu(z)$ and $U_\mu(z)$:

$$z = \cos \phi$$

$$T_\mu(z) = \cos \mu\phi = \frac{1}{2} \left[\left(-z + i\sqrt{1-z^2} \right)^\mu + \left(-z - i\sqrt{1-z^2} \right)^{-\mu} \right]$$

$$U_\mu(z) = \frac{\sin(\mu+1)\phi}{\sin \phi} = \frac{1}{\sqrt{1-z^2}} \frac{1}{2i} \left[\left(-z + i\sqrt{1-z^2} \right)^\mu + \left(-z - i\sqrt{1-z^2} \right)^{-\mu} \right]$$

We prefer to use an orthogonal basis for the product of Eqn. 5-5. So, we set:

$$\hat{T}_\mu(z) = \cotan \phi \sin \mu\phi = U_\mu(z) - T_\mu(z) \quad (5-6)$$

These functions are analytical on the complex plane, except on one cut $]-\infty, -1]$. We have the following properties:

$$(T_\mu | T_\mu) = \sin^2 \pi\mu \quad (5-7)$$

$$(\hat{T}_\mu | \hat{T}_\mu) = \tan^2 \phi \sin^2 \pi\mu \quad (5-8)$$

$$(T_\mu | \hat{T}_\mu) = 0 \quad (5-9)$$

and:

$$T_\mu(z + i0^+) - T_\mu(z - i0^-) = 4 \sin \pi\mu \sin \mu(\phi - \pi) \quad (5-10)$$

$$\hat{T}_\mu(z + i0^+) - \hat{T}_\mu(z - i0^-) = 4 \sin \pi\mu \cotan \phi \cos \mu(\phi - \pi) \quad (5-11)$$

For information, we give the asymptotics of T_μ .

- When $z \rightarrow \infty$, i.e. $x \rightarrow 0$:

$$T_\mu(z) = 2^{(\mu-1)} z^{-\mu} \left(1 - \frac{\mu}{4z^2} + \dots \right) + 2^{-(\mu+1)} z^\mu \left(1 + \frac{\mu}{4z^2} + \dots \right) \quad (5-12)$$

- When $z \rightarrow 0$, i.e. $x \rightarrow \infty$:

$$T_\mu(z) = \cos \left(\frac{\pi\mu}{2} \right) - \mu \sin \left(\frac{\pi\mu}{2} \right) z - \frac{\mu^2}{2} \cos \left(\frac{\pi\mu}{2} \right) z^2 + o(z^2) \quad (5-13)$$

$$\hat{T}_\mu(z) = \sin \left(\frac{\pi\mu}{2} \right) z - \left(\frac{\mu^2}{2} + 2\mu \right) \cos \left(\frac{\pi\mu}{2} \right) z^2 + o(z^2) \quad (5-14)$$

Case $\gamma = [0, b]$

If we perform the change of variable $z = -b/x$ sending the segment γ to $z(\gamma) =] - \infty, -1]$, we can write the general solution of Eqn. 5-4 in the form:

$$\overline{W}(x) = \frac{1}{\sin^2 \pi \mu} \left[A(x^2) \cos \mu \phi + \widehat{A}(x^2) \tan \phi \sin \mu \phi \right] \quad (5-15)$$

$$= \frac{1}{\sin^2 \pi \mu} \left[A(x^2) T_\mu(z) + \widehat{A}(x^2) \left(\frac{1}{z^2} - 1 \right) \widehat{T}_\mu(z) \right] \quad (5-16)$$

where A and $\widehat{A}$ are rational functions:

$$A(x^2) = (T_\mu | W), \quad \widehat{A}(x^2) = (\widehat{T}_\mu | W) \quad (5-17)$$

For our initial problem, A and $\widehat{A}$ are determined by the required analytical properties for $W(x)$. One can see on Eqn. 5-17 that A and $\widehat{A}$ can have singularities only at $x = \infty$, hence are polynomials. And we know the behavior of $\overline{W}$ near $x \rightarrow \infty$:

$$\overline{W}(x) = -\frac{2V'(x) - \mathbf{n}V'(-x)}{4 - \mathbf{n}^2} + o(1) \quad (5-18)$$

Thus:

$$\begin{aligned} A(x^2) &= - \left[\left(\frac{2V'(x) - \mathbf{n}V'(-x)}{4 - \mathbf{n}^2} \right) \Big| T_\mu(-b/x) \right]_+ \\ &= -\text{Even polynomial part } [V'(x)T_\mu(-b/x)] \end{aligned} \quad (5-19)$$

All the same:

$$\widehat{A}(x^2) = -\text{Even polynomial part } [V'(x)\widehat{T}_\mu(-b/x)] \quad (5-20)$$

The corresponding density is:

$$\begin{aligned} \rho(x) &= \frac{1}{2i\pi t \sin(\pi \mu)} \left[A(x^2) \sin \mu(\phi - \pi) + \widehat{A}(x^2) \tan \phi \cos \mu(\phi - \pi) \right] \\ &= \frac{\sqrt{1 - \frac{x^2}{b^2}}}{2\pi t \sin(\pi \mu)} \left[A(x^2) U_\mu(-z) + \widehat{A}(x^2) T_\mu(-z) \right] \end{aligned} \quad (5-21)$$

For example, in the fully packed case, $V'(x) = x - c/2$. When we use the asymptotics given in Eqns. 5-13 and 5-14, we find that A and $\widehat{A}$ are constants:

$$A(x^2) = b\mu \sin \left(\frac{\pi \mu}{2} \right), \quad \widehat{A}(x^2) = b \sin \left(\frac{\pi \mu}{2} \right) \quad (5-22)$$

So, we have:

$$W_1^{(0)}(x) = \frac{1}{4 \cos^2 \left(\frac{\pi \mu}{2} \right)} \left[x + \frac{b}{\sin \left(\frac{\pi \mu}{2} \right)} \left(\mu T_\mu(-b/x) + \widehat{T}_\mu(-b/x) \right) \right] \quad (5-23)$$

and the density:

$$\rho(x) = \frac{\sqrt{b^2 - x^2}}{4\pi t \cos\left(\frac{\pi\mu}{2}\right)} \left[\mu \hat{T}_\mu(-x/b) + T_\mu(-x/b) \right] \quad (5-24)$$

When b is fixed, t is the normalization such that $\int_0^b \rho = 1$.

References

- [1] J. Ambjørn, L. Chekhov, C.F. Kristjansen, Yu. Makeenko, *Matrix model calculations beyond the spherical limit*, Nucl.Phys.**B404**, pp 127-172 (1993), erratum-ibid. **B449**, p 681 (1995), [hep-th/9302014](#)
- [2] R.J. Baxter, *Exactly solved models in statistical mechanics*, Dover Publication (last edition 2007)
- [3] G. Bonnet, B. Eynard, *The Potts q -random matrix model : loop equations, critical exponents, and rational case*, Phys. Lett. **B463**, pp 273-279 (1999), [hep-th/9906130](#)
- [4] C. Berge, *Théorie des graphes et ses applications*, Dunod, 1958. Translation republished as : *The theory of graphs*, Dover Publication (2001)
- [5] M. Bergère, B. Eynard, *Universal scaling limits of matrix models, and (p, q) -Liouville gravity*, [math-ph/0909.0854](#) (2009)
- [6] J.-E. Bourgin, K. Hosomichi, *Boundary operators in the $\mathcal{O}(n)$ and RSOS matrix models*, JHEP, [hep-th/08113252](#) (2009)
- [7] E. Brézin, C. Itzykson, G. Parisi, J.-B. Zuber, *Planar diagrams*, Comm. in Math. Phys. **59**, p 35 (1978), available [here](#)
- [8] F. David, *Conformal field theories coupled to 2d gravity in the conformal gauge*, Mod. Phys. Lett. **A3**, p 1651 (1988)
- [9] J. Distler, H. Kawai, *Conformal field theory and 2d quantum gravity, or who's afraid of Joseph Liouville ?*, Nucl. Phys. **B321**, p 509 (1989)
- [10] P. Di Francesco, P. Ginsparg, J. Zinn-Justin, *2d gravity and random matrices*, Phys. Rept. **254**, pp 1-133 (1994), [hep-th/9306153v2](#)
- [11] P. Di Francesco, P. Mathieu, D. Sénéchal, *Conformal field theory*, Springer, corrected edition (1999). On the $\mathcal{O}(n)$ model : see pp 229-231.

- [12] B. Duplantier, I. Kostov, *Conformal spectra of polymers on a random surface*, Phys. Lett. Rev. **61**, p 13 (1988)
- [13] B. Durhuus, C. Kristjansen, *Phase structure of the $\mathcal{O}(\mathbf{n})$ model on a random lattice for $|\mathbf{n}| > 2$* , Nucl. Phys. **B483**, pp 535-551 (1997), [hep-th/9609008](#)
- [14] B. Eynard, *Gravité quantique bidimensionnelle et matrices aléatoires*, Thèse de doctorat (1995)
- [15] B. Eynard, *Formal matrix integrals and combinatorics of maps*, [math-ph/0611087](#) (1996)
- [16] B. Eynard, *Topological expansion for the 1-hermitian matrix model correlation functions*, JHEP 024A 0904 (2004), [hep-th/0407261](#)
- [17] B. Eynard, A. Kokotov, D. Korotkin, *Genus one contribution to free energy in hermitian two-matrix model*, Nucl. Phys. **B694**, pp 443-472 (2004), [hep-th/0403072](#)
- [18] B. Eynard, C. Kristjansen, *Exact solution of the $\mathcal{O}(\mathbf{n})$ model on a random lattice*, Nucl. Phys. **B455**, pp 577-618 (1995) [hep-th/9506193](#)
- [19] B. Eynard, C. Kristjansen, *More on the exact solution of the $\mathcal{O}(\mathbf{n})$ model on a random lattice and an investigation of the case $|\mathbf{n}| > 2$* , Nucl. Phys. **B466**, pp 463-487 (1996), [hep-th/9512052](#)
- [20] B. Eynard, N. Orantin, *Invariants of algebraic curves and topological expansion*, Communications in Number Theory and Physics, Vol 1, Number 2, pp 347-452 (2007), [math-ph/0702045](#)
- [21] B. Eynard, N. Orantin, *Topological expansion and boundary conditions*, JHEP 0806 037 (2008), [hep-th/0710.0223](#)
- [22] B. Eynard, N. Orantin, *Geometrical interpretation of the topological recursion, and integrable string theory*, [math-ph/0911.5096](#) (2009)
- [23] B. Eynard, J. Zinn-Justin, *The $\mathcal{O}(\mathbf{n})$ model on a random surface: critical points and large order behavior*, Nucl. Phys. **B386**, pp 558-591 (1992) [hep-th/9204082](#)
- [24] M. Gaudin, I. Kostov, *$\mathcal{O}(\mathbf{n})$ on a fluctuating planar lattice. Some exact results*, Phys. Lett. **B220**, p 200 (1989)
- [25] Z.X. Wang, D.R. Guo, *Special functions*, ed. World Scientific (1989)

- [26] V. Kazakov, I. Kostov, *Loop gas model for open strings*, Nucl. Phys. **B386**, p 520 (1992), [hep-th/9205059](#)
- [27] V. G. Knizhnik, A. M. Polyakov, A. B. Zamolodchikov, *Fractal structure of 2d quantum gravity*, Mod. Phys. Lett. **A3**, p 819 (1988)
- [28] J. M. Kosterlitz, *The critical properties of the two dimensional $x - y$ model*, J. Phys. **C7**, p 1046 (1974)
- [29] I. Kostov, M.L. Mehta, *Random surfaces of arbitrary genus: exact results for $D = 0$, and -2 dimensions*, Phys. Lett. **B189**, pp 118-124 (1987)
- [30] I. Kostov, *$\mathcal{O}(n)$ vector model on a planar random lattice: spectrum of anomalous dimensions*, Mod. Phys. Lett. **A4**, p 217 (1989)
- [31] I. Kostov, *Loop amplitude for nonrational string theories*, Phys. Lett. **B266**, p 312 (1991)
- [32] I. Kostov, M. Staudacher, *Multicritical phases of the $\mathcal{O}(n)$ model on a random lattice*, Nucl. Phys. **B384**, pp 459-483 (1992), [hep-th/9203030](#)
- [33] I. Kostov, *Gauge invariant matrix models for the $\hat{A} - \hat{D} - \hat{E}$ closed strings*, Phys. Lett. **B297** pp 74-81 (1992), [hep-th/9208053](#)
- [34] I. Kostov, *Boundary loop models and 2D quantum gravity*, J. Stat. Mech. 07-08, pp 8-23 (2007) [hep-th/0703221](#)
- [35] J. L. Jacobsen, H. Saleur, *Conformal boundary loop models*, Nucl. Phys. **B788**, pp 137-166 (2008), [math-ph/0611078](#)
- [36] A.A. Migdal, Phys. Rep. **102**, p 199 (1983)
- [37] G.W. Moore, N. Seiberg, M. Staudacher, *From loops to states in 2d quantum gravity*, Nucl. Phys. **B362**, pp 665-709 (1991)
- [38] Y. Nakayama, *Liouville Field Theory - A decade after the revolution*, Int. J. Mod. Phys. **A19**, pp 2771-2930 (2004), [hep-th/0402009](#)
- [39] F.G. Tricomi, *Integral Equations*, Pure Appl. Math. V, Interscience, London (1957)
- [40] W.T. Tutte, *A census of planar triangulations*, Can. J. Math. **14**, pp 21-38 (1962)
- [41] W.T. Tutte, *A census of planar maps*, Can. J. Math. **15**, pp 249-271 (1963)
- [42] P. Zinn-Justin, *The asymmetric ABAB model*, Europhys. Lett. **64** pp 737-742 (2003), [hep-th/0308132](#)